

Recombinant Human EIF3G 293 Cell Lysate

Cat. No. EIF3G-6660HCL Lot. No. (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for eukaryotic translation initiation factor 3, subunit G (EIF3G) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	EIF3G eukaryotic translation initiation factor 3, subunit G [Homo sapiens]
Official Symbol	EIF3G
Synonyms	EIF3G; eukaryotic translation initiation factor 3, subunit G; EIF3S4, eukaryotic translation initiation factor 3, subunit 4 delta, 44kDa; eukaryotic translation initiation factor 3 subunit G; eIF3 delta; eIF3 p44; eIF3g; eIF3 p42; eIF-3-delta; eIF-3 RNA-binding subunit; eukaryotic translation initiation factor 3 subunit 4; eukaryotic translation initiation factor 3 subunit p42; eukaryotic translation initiation factor 3 RNA-binding subunit; eukaryotic translation initiation factor 3, subunit 4 delta, 44kDa; eukaryotic translation initiation factor 3, subunit 4 (delta, 44kD); EIF3S4; EIF3-P42; eIF3-p44; eIF3-delta;
Gene ID	8666
mRNA Refseq	NM_003755
Protein Refseq	NP_003746
MIM	603913
UniProt ID	O75821
Chromosome Location	19p13.2
Pathway	Activation of the mRNA upon binding of the cap-binding complex and eIFs, and

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

subsequent binding to 43S, organism-specific biosystem; Cap-dependent Translation Initiation, organism-specific biosystem; Eukaryotic Translation Initiation, organism-specific biosystem; Formation of a pool of free 40S subunits, organism-specific biosystem; Formation of the ternary complex, and subsequently, the 43S complex, organism-specific biosystem; GTP hydrolysis and joining of the 60S ribosomal subunit, organism-specific biosystem; Gene Expression, organism-specific biosystem;

Function

RNA binding; nucleotide binding; protein binding; contributes_to translation initiation factor activity; contributes_to translation initiation factor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA