

Recombinant Human EIF4A2 293 Cell Lysate

Cat. No. EIF4A2-6653HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for eukaryotic translation initiation factor 4A2 (EIF4A2) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	EIF4A2 eukaryotic translation initiation factor 4A2 [Homo sapiens]
Official Symbol	EIF4A2
Synonyms	EIF4A2; eukaryotic translation initiation factor 4A2; EIF4F, eukaryotic translation initiation factor 4A, isoform 2; eukaryotic initiation factor 4A-II; BM 010; DDX2B; EIF4A; ATP-dependent RNA helicase eIF4A-2; eukaryotic translation initiation factor 4A, isoform 2; EIF4F; BM-010; eIF4A-II; eIF-4A-II;
Gene ID	1974
mRNA Refseq	NM_001967
Protein Refseq	NP_001958
MIM	601102
UniProt ID	Q14240
Chromosome Location	3q28
Pathway	Activation of the mRNA upon binding of the cap-binding complex and eIFs, and subsequent binding to 43S, organism-specific biosystem; Antiviral mechanism by IFN-stimulated genes, organism-specific biosystem; Cap-dependent Translation Initiation, organism-specific biosystem; Cytokine Signaling in Immune system, organism-specific biosystem; Deadenylation of mRNA, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA



Deadenylation-dependent mRNA decay, organism-specific biosystem; Eukaryotic Translation Initiation, organism-specific biosystem;

Function

ATP binding; ATP-dependent helicase activity; RNA binding; helicase activity; hydrolase activity; nucleic acid binding; nucleotide binding; protein binding; translation initiation factor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA