

Recombinant Human EIF5A 293 Cell Lysate

Cat. No. EIF5A-6641HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for eukaryotic translation initiation factor 5A (EIF5A), transcript variant B is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	EIF5A eukaryotic translation initiation factor 5A [Homo sapiens]
Official Symbol	EIF5A
Synonyms	EIF5A; eukaryotic translation initiation factor 5A; eukaryotic translation initiation factor 5A-1; EIF 5A; EIF5A1; MGC99547; MGC104255; eIF-4D; eIF-5A1; eIF-5A-1; rev-binding factor; eukaryotic initiation factor 5A; EIF-5A; eIF5A1;
Gene ID	1984
mRNA Refseq	NM_001143760
Protein Refseq	NP_001137232
MIM	600187
UniProt ID	P63241
Chromosome Location	17p13-p12
Pathway	Hypusine synthesis from eIF5A-lysine, organism-specific biosystem; Metabolism of proteins, organism-specific biosystem; PTM: gamma carboxylation, hypusine formation and arylsulfatase activation, organism-specific biosystem; Post-translational protein modification, organism-specific biosystem; Translation Factors, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

RNA binding; U6 snRNA binding; protein N-terminus binding; protein binding; ribosome binding; translation elongation factor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA