

Active Recombinant Human EPO protein, His-tagged

Cat. No. EPO-451H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human EPO protein(P01588)(28~193 aa), fused with N-terminal His tag, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	28-193 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	<p>Erythropoietin (EPO), also known as hematopoietin or hemopoietin, is a glycoprotein cytokine secreted by the kidney in response to cellular hypoxia. Erythropoietin is an essential hormone for red blood cell production. Its primary effect on red blood cell progenitors and precursors (which are found in the bone marrow in humans) by promoting their survival through protecting these cells from apoptosis, or cell death. EPO is the primary erythropoietic factor that cooperates with various other growth factors involved in the development of erythroid lineage from multipotent progenitors. Besides, Erythropoietin Receptor (EPOR) has been identified as an interactor of EPO, thus a binding ELISA assay was conducted to detect the interaction of recombinant human EPO and recombinant human EPOR. Briefly, EPO were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100uL were then transferred to EPOR-coated microtiter wells and incubated for 2h at 37°C. Wells were</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

washed with PBST and incubated for 1h with anti-EPO pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µ stop solution to the wells and read at 450nm immediately. The binding activity of EPO and EPOR was shown, and this effect was in a dose dependent manner.

Molecular Mass	20.0kDa
Endotoxin	<1.0EU per 1µg (determined by the LAL method).
Purity	> 95% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 10mM PBS (pH7.6) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	EPO erythropoietin [Homo sapiens]
Official Symbol	EPO
Synonyms	EPO; erythropoietin; EP; epoetin; MVCD2; MGC138142;
Gene ID	2056
mRNA Refseq	NM_000799

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq NP_000790


MIM 133170

UniProt ID P01588

SDS-PAGE



Activity



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA