

Recombinant Human ERBB2 cell lysate

Cat. No. ERBB2-2658HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human ErbB2 (HER2) Protein, was expressed in Human Cells (whole cell lysate provided in 1X Sample Buffer; note: transfected cell lysate positive control).
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min.3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles.Notes:The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

ERBB2 v-erb-b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog (avian) [Homo sapiens]

Official Symbol

ERBB2

Synonyms

ERBB2; v-erb-b2 erythroblastic leukemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog (avian); NGL, v erb b2 avian erythroblastic leukemia viral oncogene homolog 2 (neuro/glioblastoma derived oncogene homolog); receptor tyrosine-protein kinase erbB-2; CD340; HER 2; HER2; NEU; herstatin; p185erbB2; proto-oncogene Neu; c-erb B2/neu protein; proto-oncogene c-ErbB-2; metastatic lymph node gene 19 protein; tyrosine kinase-type cell surface receptor HER2; neuroblastoma/glioblastoma derived oncogene homolog; v-erb-b2 avian erythroblastic leukemia viral oncogene homolog 2 (neuro/glioblastoma derived oncogene homolog); NGL; TKR1; HER-2; MLN 19; HER-2/neu;

Gene ID

2064

mRNA Refseq

NM_001005862

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq	NP_001005862
MIM	164870
UniProt ID	P04626
Chromosome Location	17q11.2-q12
Pathway	Adherens junction, organism-specific biosystem; Adherens junction, conserved biosystem; Alpha6-Beta4 Integrin Signaling Pathway, organism-specific biosystem; Axon guidance, organism-specific biosystem; Bladder cancer, organism-specific biosystem; Bladder cancer, conserved biosystem; Calcium signaling pathway, organism-specific biosystem;
Function	ATP binding; ErbB-3 class receptor binding; Hsp90 protein binding; RNA polymerase I core binding; epidermal growth factor-activated receptor activity; glycoprotein binding; contributes_to growth factor binding; identical protein binding; nucleotide binding; protein C-terminus binding; protein binding; protein dimerization activity; protein heterodimerization activity; protein heterodimerization activity; protein heterodimerization activity; protein phosphatase binding; protein tyrosine kinase activity; protein tyrosine kinase activity; receptor activity; receptor signaling protein tyrosine kinase activity; transmembrane receptor protein tyrosine kinase activity; transmembrane signaling receptor activity; ubiquitin protein ligase binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA