

Recombinant Human ERCC3 protein, His & T7-tagged

Cat. No. ERCC3-1692H Lot. No. (See product label)

SPECIFICATION

| | |
|-------------------------|--|
| Product Overview | Recombinant Human ERCC3 aa. (Arg542~Ala702 (Accession # P19447)) fused with N-terminal His & T7 tag was produced in E. coli cells. |
| Species | Human |
| Source | E.coli |
| ProteinLength | Arg542~Ala702 |
| Description | This gene encodes an ATP-dependent DNA helicase that functions in nucleotide excision repair. The encoded protein is a subunit of basal transcription factor 2 (TFIIH) and, therefore, also functions in class II transcription. Mutations in this gene are associated with Xeroderma pigmentosum B, Cockayne's syndrome, and trichothiodystrophy. Alternative splicing results in multiple transcript variants. |
| Form | Freeze-dried powder |
| Molecular Mass | Predicted Molecular Mass: 22.3kDa |
| Endotoxin | <1.0EU per 1ug (determined by the LAL method) |
| Purity | >95% |
| Applications | SDS-PAGE; WB; ELISA; IP. |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

| | |
|--------------------------|---|
| Stability | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Storage | Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months. |
| Storage buffer | Supplied as lyophilized form in 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5% trehalose, and preservative. |
| Reconstitution | Reconstitute in sterile ddH ₂ O. |
| Isoelectric Point | 9.5 |

GENE INFORMATION

| | |
|------------------------|---|
| Gene Name | ERCC3 ERCC excision repair 3, TFIIH core complex helicase subunit [Homo sapiens (human)] |
| Official Symbol | ERCC3 |
| Synonyms | ERCC3; ERCC excision repair 3, TFIIH core complex helicase subunit; XPB; BTF2; TTD2; GTF2H; RAD25; TFIIH; TFIIH basal transcription factor complex helicase XPB subunit; BTF2 p89; DNA excision repair protein ERCC-3; DNA repair helicase; DNA repair protein complementing XP-B cells; TFIIH 89 kDa subunit; TFIIH basal transcription factor complex 89 kDa subunit; TFIIH p89; basic transcription factor 2 89 kDa subunit; excision repair cross-complementation group 3; excision repair cross-complementing rodent repair deficiency, complementation group 3; xeroderma pigmentosum group B-complementing protein; xeroderma pigmentosum, complementation group B |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


Gene ID 2071

mRNA Refseq NM_000122.1

Protein Refseq NP_000113.1

UniProt ID P19447

SDS-PAGE



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA