

## Active Recombinant Mouse Epcam protein, Fc-tagged

Cat. No. Epcam-191M Lot. No. (See product label)

### SPECIFICATION

**Product Overview** Recombinant Mouse Epcam(Gln24 & Arg81-Thr266) fused with Fc region of Mouse IgG2a at C-terminal was expressed in NS0.

**Species** Mouse

**Source** Mammalian Cells

**ProteinLength** 81-266 a.a.

#### Description

Epithelial Cellular Adhesion Molecule (EpCAM), also known as KS1/4, gp40, GA733-2, 17-1A, CD326 and TROP-1, is a 40 kDa transmembrane glycoprotein composed of a 243 amino acid (aa) extracellular domain with two epidermal-growth-factor-like (EGF-like) repeats, a 23 aa transmembrane domain, and a 26 aa cytoplasmic domain. Human and mouse EpCAM share 82% aa sequence identity. In mouse, EpCAM also shares 51% aa sequence homology with Trop-2. EpCAM is detected in basolateral cell membranes of all simple epithelia and expression has been found to increase in epithelia tissues during fetal development. EpCAM has been shown to function as a homophilic Ca<sup>2+</sup> independent adhesion molecule, but it does not structurally resemble any of the four major families of cell adhesion molecules. Homophilic adhesion via EpCAM requires the interaction of both EGF-like repeats, with the first EGF-like repeat mediating reciprocal interaction between EpCAM molecules on opposing cells, while the second repeat is involved in lateral interaction of EpCAM. EpCAM has been identified as a surface marker for pluripotent embryonic stem cells and is

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	strongly associated with the maintenance of the undifferentiated state of ESCs. EpCAM is highly expressed on the surface of epithelial cancer cells including colon, stomach, lung, pancreas, prostate and breast.
<b>Predicted N Terminal</b>	Arg81 detected. Gln24 inferred from enzymatic pyroglutamate treatment revealing Arg25
<b>Form</b>	Lyophilized from a 0.2 µm filtered solution in PBS.
<b>Bio-activity</b>	Measured by the ability of the immobilized protein to support the adhesion of the L Cells mouse fibroblast cell line. When 5 x 10 <sup>4</sup> cells/well are added to Recombinant Mouse EpCAM/TROP-1 and Human Fibronectin coated plates, cell adhesion is enhanced in a dose dependent manner. The ED50 for this effect is typically 0.4-2.4 µg/mL.
<b>Molecular Mass</b>	Predicted Molecular Mass: 55 kDa SDS-PAGE: 58-75 kDa and 8 kDa
<b>Endotoxin</b>	<0.1 EU per 1 µg of the protein by the LAL method.
<b>Purity</b>	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
<b>Storage</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
<b>Reconstitution</b>	Reconstitute at 500 µg/mL in PBS.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

## GENE INFORMATION

**Gene Name** Epcam epithelial cell adhesion molecule [ Mus musculus ]

**Official Symbol** Epcam

**Synonyms** EPCAM; epithelial cell adhesion molecule; mEGP314; protein 289A; Trop-1 protein; lymphocyte antigen 74; epithelial glycoprotein 314; panepithelial glycoprotein 314; tumor-associated calcium signal transducer 1; EGP; Ly74; gp40; CD326; EGP-2; TROP1; Egp314; Ep-CAM; Tacsd1; GA733-2; Tacstd1;

**Gene ID** 17075

**mRNA Refseq** NM\_008532

**Protein Refseq** NP\_032558

**UniProt ID** Q99JW5

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA