

Active Recombinant Mouse Epcam Protein, Fc-tagged, Alexa Fluor 488 conjugated

Cat. No. Epcam-191MAF488 **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Alexa Fluor 488 conjugated recombinant Mouse Epcam(Gln24 & Arg81-Thr266) fused with Fc region of Mouse IgG2a at C-terminal was expressed in NS0.

Species

Mouse

Source

Mammalian Cells

ProteinLength

81-266 a.a.

Description

Epithelial Cellular Adhesion Molecule (EpCAM), also known as KS1/4, gp40, GA733-2, 17-1A, CD326 and TROP-1, is a 40 kDa transmembrane glycoprotein composed of a 243 amino acid (aa) extracellular domain with two epidermal-growth-factor-like (EGF-like) repeats, a 23 aa transmembrane domain, and a 26 aa cytoplasmic domain. Human and mouse EpCAM share 82% aa sequence identity. In mouse, EpCAM also shares 51% aa sequence homology with Trop-2. EpCAM is detected in basolateral cell membranes of all simple epithelia and expression has been found to increase in epithelia tissues during fetal development. EpCAM has been shown to function as a homophilic Ca²⁺ independent adhesion molecule, but it does not structurally resemble any of the four major families of cell adhesion molecules. Homophilic adhesion via EpCAM requires the interaction of both EGF-like repeats, with the first EGF-like repeat mediating reciprocal interaction between EpCAM molecules on opposing cells, while the second repeat is involved in lateral interaction of EpCAM. EpCAM has

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

been identified as a surface marker for pluripotent embryonic stem cells and is strongly associated with the maintenance of the undifferentiated state of ESCs. EpCAM is highly expressed on the surface of epithelial cancer cells including colon, stomach, lung, pancreas, prostate and breast.

Form

Lyophilized

Bio-activity

Measured by the ability of the immobilized protein to support the adhesion of the L Cells mouse fibroblast cell line. When 5×10^4 cells/well are added to Recombinant Mouse EpCAM/TROP-1 and Human Fibronectin coated plates, cell adhesion is enhanced in a dose dependent manner. The ED50 for this effect is typically 0.4-2.4 $\mu\text{g/mL}$.

Molecular Mass

Predicted Molecular Mass: 55 kDa
SDS-PAGE: 58-75 kDa and 8 kDa

N-terminal Sequence Analysis

Arg 81 detected. Gln 24 inferred from enzymatic pyroglutamate treatment revealing Arg 25.

Endotoxin

< 0.1 EU/ μg of the protein by the LAL method.

Purity

> 95 % by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

Characteristic

Disulfide-linked homodimer
Labeled with Alexa Fluor 488 via amines
Excitation Wavelength: 488 nm
Emission Wavelength: 515-545 nm

Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



12 months from date of receipt, -20 to -70 centigrade as supplied.
1 month, 2 to 8 centigrade under sterile conditions after reconstitution.
3 months, -20 to -70 centigrade under sterile conditions after reconstitution.

Concentration	500 µg/mL
Storage Buffer	Lyophilized from a 0.2 µm filtered solution in PBS.
Reconstitution	Reconstitute at 500 µg/mL in PBS.
Conjugation	Alexa Fluor 488

GENE INFORMATION

Gene Name	Epcam epithelial cell adhesion molecule [Mus musculus]
Official Symbol	Epcam
Synonyms	EPCAM; epithelial cell adhesion molecule; mEGP314; protein 289A; Trop-1 protein; lymphocyte antigen 74; epithelial glycoprotein 314; panepithelial glycoprotein 314; tumor-associated calcium signal transducer 1; EGP; Ly74; gp40; CD326; EGP-2; TROP1; Egp314; Ep-CAM; Tacsd1; GA733-2; Tacstd1;
Gene ID	17075
mRNA Refseq	NM_008532
Protein Refseq	NP_032558
UniProt ID	Q99JW5