

## Recombinant Human FGF18 cell lysate

**Cat. No.** FGF18-1660HCL    **Lot. No.** (See product label)

### SPECIFICATION

<b>Product Overview</b>	Human FGF18 / FGF-18 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
<b>Species</b>	Human
<b>Source</b>	Human Cells
<b>Preparation method</b>	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
<b>Lysis buffer</b>	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
<b>Quality control Testing</b>	12.5% SDS-PAGE Stained with Coomassie Blue
<b>Recommended Usage</b>	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

**Stability**

Samples are stable for up to twelve months from date of receipt at -80°C

**Storage Buffer**

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

**Storage Instruction**

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

## GENE INFORMATION

**Gene Name**

FGF18 fibroblast growth factor 18 [ Homo sapiens ]

**Official Symbol**

FGF18

**Synonyms**

FGF18; fibroblast growth factor 18; FGF 18; ZFGF5; FGF-18;

**Gene ID**

8817

**mRNA Refseq**

NM\_003862

**Protein Refseq**

NP\_003853

**MIM**


603726

**UniProt ID**

O76093

**Chromosome  
Location**

5q34

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

**Pathway**

Downstream signaling of activated FGFR, organism-specific biosystem;  
Endochondral Ossification, organism-specific biosystem; FGFR ligand binding and activation, organism-specific biosystem; FGFR2 ligand binding and activation, organism-specific biosystem; FGFR2c ligand binding and activation, organism-specific biosystem; FGFR3 ligand binding and activation, organism-specific biosystem; FGFR3b ligand binding and activation, organism-specific biosystem;

**Function**

growth factor activity; protein tyrosine kinase activity; type 1 fibroblast growth factor receptor binding; type 2 fibroblast growth factor receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA