

Active Recombinant Human FGF2 Protein (Formulation I- Animal Free-Lyophilized, 134-288, 154 amino acid)

Cat. No. FGF2-20H Lot. No. (See product label)

SPECIFICATION

Product Overview

Recombinant Human FGF2 Protein (Formulation I- Animal Free-Lyophilized, 134-288, 154 amino acid) without tag was expressed in E. coli.

Species

Human

Source

E.coli

ProteinLength

134-288, 154 amino acid

Description

The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members bind heparin and possess broad mitogenic and angiogenic activities. This protein has been implicated in diverse biological processes, such as limb and nervous system development, wound healing, and tumor growth. The mRNA for this gene contains multiple polyadenylation sites, and is alternatively translated from non-AUG (CUG) and AUG initiation codons, resulting in five different isoforms with distinct properties. The CUG-initiated isoforms are localized in the nucleus and are responsible for the intracrine effect, whereas, the AUG-initiated form is mostly cytosolic and is responsible for the paracrine and autocrine effects of this FGF.

Form

Lyophilized

Bio-activity

ED50= 0.1-1.0 ng/mL as determined by the dose dependent proliferation of NIH 3T3 cells.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	17.3 kDa
AA Sequence	AAGSITTLPALPEDGGSGAFPPGHFKDPKRLYCKNGGFFLRIHPDGRVDGVREKSD PHIKLQLQAEERGVSIVKGVCANRYLAMKEDGRLLASKCVTDECFERLESNNYNT YRSRKYTSWYVALKRTGQYKLGSKTGPQKAILFLPMSAKS
Endotoxin	< 0.01 ng/μg cytokine as determined by the LAL assay
Purity	> 98% by SDS-PAGE
Usage	Useful for cell culture and for the study of signaling pathways.
Storage	Upon receipt, it should be stored immediately at -20 to -80 centigrade. Lyophilized bFGF is stable at -20 to -80 centigrade for up to 16 months.
Storage Buffer	Sterile filtered through a 0.22 micron filter and lyophilized from Tris buffer (10 mM Tris pH7.5, 150 mM NaCl).
Reconstitution	Centrifuge vial before opening. It is recommended to reconstitute bFGF in sterile 10 mM Tris-HCl, pH 7.5 to yield a stock solution of 0.1 mg/mL of bFGF. It is stable for up to 6 months when stored at -20 centigrade and up to 12 months when stored at -80 centigrade. Multiple freeze/thaw cycles will result in significant loss of activity.
Shipping	The product is shipped at ambient temperature with ice bags.

GENE INFORMATION

Gene Name	FGF2 fibroblast growth factor 2 [Homo sapiens (human)]
Official Symbol	FGF2


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Synonyms	FGF2; fibroblast growth factor 2; BFGF; FGFB; FGF-2; HBGF-2; fibroblast growth factor 2; basic fibroblast growth factor bFGF; fibroblast growth factor 2 (basic); heparin-binding growth factor 2; prostatropin
Gene ID	2247
mRNA Refseq	NM_002006
Protein Refseq	NP_001997
MIM	134920
UniProt ID	P09038

SDS-PAGE




The image shows an SDS-PAGE gel with a single prominent band labeled 'bFGF' at approximately 20 kDa. To the right of the gel, molecular weight markers are indicated at 160kDa, 110kDa, 80kDa, 60kDa, 50kDa, 40kDa, 30kDa, 20kDa, 15kDa, and 10kDa.

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

Activity



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA