

Recombinant Human FGF2 cell lysate

Cat. No. FGF2-638HCL Lot. No. (See product label)

SPECIFICATION

Product Overview	Human VEGF121 / VEGF-A derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

FGF2 fibroblast growth factor 2 (basic) [Homo sapiens]

Official Symbol

FGF2

Synonyms

FGF2; fibroblast growth factor 2 (basic); FGFB; fibroblast growth factor 2; prostatropin; heparin-binding growth factor 2; basic fibroblast growth factor bFGF; BFGF; FGF-2; HBGF-2;

Gene ID

2247

mRNA Refseq

NM_002006

Protein Refseq

NP_001997

MIM


134920

UniProt ID

P09038

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	4q26
Pathway	Angiopoietin receptor Tie2-mediated signaling, organism-specific biosystem; Downstream signaling of activated FGFR, organism-specific biosystem; Endochondral Ossification, organism-specific biosystem; FGFR ligand binding and activation, organism-specific biosystem; FGFR1 ligand binding and activation, organism-specific biosystem; FGFR1b ligand binding and activation, organism-specific biosystem; FGFR1c ligand binding and activation, organism-specific biosystem;
Function	chemoattractant activity; cytokine activity; fibroblast growth factor binding; fibroblast growth factor receptor binding; growth factor activity; heparin binding; ligand-dependent nuclear receptor transcription coactivator activity; protein binding; protein tyrosine kinase activity; voltage-gated calcium channel activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA