

Recombinant Active Human FGF4 Protein, His-tagged(C-ter)

Cat. No. FGF4-86H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Active Human FGF4 Protein with His tag (C-ter) was expressed in E. coli.
Species	Human
Source	E.coli
Description	The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities and are involved in a variety of biological processes including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This gene was identified by its oncogenic transforming activity. This gene and FGF3, another oncogenic growth factor, are located closely on chromosome 11. Co-amplification of both genes was found in various kinds of human tumors. Studies on the mouse homolog suggested a function in bone morphogenesis and limb development through the sonic hedgehog (SHH) signaling pathway. [provided by RefSeq, Jul 2008]
Form	Powder
Bio-activity	Determined by its ability to induce 3T3 cells proliferation. The ED50 for this effect is < 2.5 ng/mL. The specific activity of recombinant human FGF-4 is > 4 x 10 ⁵ IU/mg.
AA Sequence	MGLECDGKVNICCKKQFFVFSKDIGWNDWIIAPSGYHANYCEGECPSHIAGTSGSSL SFHSTVINHYRMRGHSPFANLKSCCVPTKLRPMSMLYDDGQNIKKDIQNMIVEEC GCS

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	Endotoxin level is less than 0.1 EU/μg of the protein, as determined by the LAL test.
Purity	> 95% (by SDS-PAGE)
Applications	SDS-PAGE
Notes	For laboratory research only, not for drug, diagnostic or other use.
Storage	Lyophilized protein should be stored at -20°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Storage Buffer	PBS (pH 8.0) and 0.1% Sarkosyl.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.

GENE INFORMATION


Gene Name	FGF4 fibroblast growth factor 4 [Homo sapiens]
Official Symbol	FGF4
Synonyms	FGF4; fibroblast growth factor 4; heparin secretory transforming protein 1 , HSTF1; HBGF 4; HST; HST 1; human stomach cancer; transforming factor from FGF related oncogene; K FGF; kaposi sarcoma oncogene; KFGF; transforming protein KS3; FGF-4; HSTF-1; oncogene HST; heparin-binding growth factor 4; heparin secretory transforming protein 1; heparin secretory-transforming protein 1; fibroblast growth factor 4 splice isoform; human stomach cancer, transforming factor from FGF-related

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

oncogene; HST-1; HSTF1; K-FGF; HBGF-4;

Gene ID 2249**mRNA Refseq** NM_002007**Protein Refseq** NP_001998**MIM** 164980**UniProt ID** P08620**SDS-PAGE**Human FGF4
recombinant protein Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA