

## Recombinant Mouse FGFR1 cell lysate

**Cat. No.** FGFR1-2133MCL    **Lot. No.** (See product label)

### SPECIFICATION

<b>Product Overview</b>	Mouse FGFR1 / CD331 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
<b>Species</b>	Mouse
<b>Source</b>	Human Cells
<b>Preparation method</b>	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
<b>Lysis buffer</b>	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
<b>Quality control Testing</b>	12.5% SDS-PAGE Stained with Coomassie Blue
<b>Recommended Usage</b>	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

**Stability**

Samples are stable for up to twelve months from date of receipt at -80°C

**Storage Buffer**

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

**Storage Instruction**

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

## GENE INFORMATION

**Gene Name**

[Fgfr1 fibroblast growth factor receptor 1 \[ Mus musculus \]](#)

**Official Symbol**

FGFR1

**Synonyms**

FGFR1; fibroblast growth factor receptor 1; MFR; c-fgr; bFGF-R-1; proto-oncogene c-Fgr; basic fibroblast growth factor receptor 1; FLG; Eask; Hspy; Flt-2; Fgfr-1; AW208770;

**Gene ID**

[14182](#)

**mRNA Refseq**

[NM\\_001079908](#)

**Protein Refseq**

[NP\\_001073377](#)

**Pathway**

Adherens junction, organism-specific biosystem; Adherens junction, conserved biosystem; Axon guidance, organism-specific biosystem; Developmental Biology, organism-specific biosystem; Downstream signaling of activated FGFR, organism-specific biosystem; ESC Pluripotency Pathways, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endochondral Ossification, organism-specific biosystem;

**Function**

ATP binding; cell adhesion molecule binding; fibroblast growth factor 1 binding; fibroblast growth factor binding; fibroblast growth factor binding; fibroblast growth factor-activated receptor activity; fibroblast growth factor-activated receptor activity; glycoprotein binding; heparin binding; kinase activity; nucleotide binding; protein binding; protein complex binding; protein homodimerization activity; protein kinase activity; protein tyrosine kinase activity; receptor activity; transferase activity; transferase activity, transferring phosphorus-containing groups; transmembrane receptor protein tyrosine kinase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA