

# Recombinant Human FGFR1

Cat. No. FGFR1-28890TH Lot. No. (See product label)

## SPECIFICATION

**Product Overview**

Recombinant fusion protein: Soluble FGFR1a Fc Chimera of Recombinant Human Soluble FGFR1a fused via a Xa cleavage site with the Fc part of human IgG1. Produced by baculovirus in insect cells. A heterodimeric, glycosylated, polypeptide chain having a molecule

**Species** Human

**Description**

The protein encoded by this gene is a member of the fibroblast growth factor receptor (FGFR) family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein consists of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member binds both acidic and basic fibroblast growth factors and is involved in limb induction. Mutations in this gene have been associated with Pfeiffer syndrome, Jackson-Weiss syndrome, Antley-Bixler syndrome, osteoglophonic dysplasia, and autosomal dominant Kallmann syndrome 2. Chromosomal aberrations involving this gene are associated with stem cell myeloproliferative disorder and stem cell leukemia lymphoma syndrome. Alternatively spliced variants which encode different protein isoforms have been described; however, not all variants have been fully characterized.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Tissue specificity</b>	Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.
<b>Form</b>	Lyophilised:Reconstitute in sterile PBS not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.
<b>Purity</b>	>90% by SDS-PAGE
<b>Storage buffer</b>	Preservative: NoneNo additives
<b>Storage</b>	Aliquot and store at -80°C. Avoid repeated freeze / thaw cycles.
<b>Sequence Similarities</b>	Belongs to the protein kinase superfamily. Tyr protein kinase family. Fibroblast growth factor receptor subfamily.Contains 3 Ig-like C2-type (immunoglobulin-like) domains.Contains 1 protein kinase domain.

## GENE INFORMATION

<b>Gene Name</b>	FGFR1 fibroblast growth factor receptor 1 [ Homo sapiens ]
<b>Official Symbol</b>	FGFR1
<b>Synonyms</b>	FGFR1; fibroblast growth factor receptor 1; FLT2, fms related tyrosine kinase 2 , KAL2; BFGFR; CD331; CEK; FLG; H2; H3; H4; H5; N SAM; Pfeiffer syndrome;
<b>Gene ID</b>	2260
<b>mRNA Refseq</b>	NM_001174063
<b>Protein Refseq</b>	NP_001167534

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>MIM</b>	136350
<b>Uniprot ID</b>	P11362
<b>Chromosome Location</b>	8p12
<b>Pathway</b>	Adherens junction, organism-specific biosystem; Adherens junction, conserved biosystem; Axon guidance, organism-specific biosystem; Developmental Biology, organism-specific biosystem; Downstream signaling of activated FGFR, organism-specific biosystem;
<b>Function</b>	ATP binding; fibroblast growth factor 1 binding; fibroblast growth factor binding; fibroblast growth factor-activated receptor activity; fibroblast growth factor-activated receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA