

Recombinant Human Fms-related Tyrosine Kinase 1, 4 Domains

Cat. No. FLT1-79H **Lot. No.** (See product label)

SPECIFICATION

| | |
|-------------------------|---|
| Product Overview | Recombinant Human Fms-related Tyrosine Kinase 1 D1-4 produced in baculovirus is monomeric, glycosylated, polypeptide containing 457 amino acids and having a molecular mass of 55 kDa. The soluble receptor protein contains only the first 4 extracellular domains, which contain all the information necessary for binding of VEGF. The VEGFR1 is purified by proprietary chromatographic techniques. |
| Species | Human |
| Source | Insect Cells |
| Description | <p>Endothelial cells express three different vascular endothelial growth factor (VEGF) receptors, belonging to the family of receptor tyrosine kinases (RTKs). They are named VEGFR-1 (Flt-1), VEGFR-2 (KDR/Flk-1), VEGFR-3 (Flt-4). Their expression is almost exclusively restricted to endothelial cells, but VEGFR-1 can also be found on monocytes, dendritic cells and on trophoblast cells. The flt-1 gene was first described in 1990. The receptor contains seven immunoglobulin-like extracellular domains, a single transmembrane region and an intracellular split tyrosine kinase domain. Compared to VEGFR-2 the Flt-1 receptor has a higher affinity for VEGF but a weaker signaling activity. VEGFR-1 thus leads not to proliferation of endothelial cells, but mediates signals for differentiation. Interestingly a naturally occurring soluble variant of VEGFR-1 (sVEGFR-1) was found in HUVE supernatants in 1996, which is generated by alternative splicing of the flt-1 mRNA. The biological functions of sVEGFR-1 still are not clear, but it seems to be an endogenous regulator of angiogenesis, binding</p> |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

| | |
|----------------------------|--|
| | VEGF with the same affinity as the full-length receptor. |
| Physical Appearance | Sterile Filtered White lyophilized (freeze-dried) powder. |
| Purity | Greater than 90.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.GE. |
| Formulation | VEGFR-1 D1-4 was lyophilized from a concentrated (1mg/ml) sterile solution containing no additives. |
| Solubility | It is recommended to reconstitute the lyophilized sVEGFR-1 D4 in sterile water not less than 100 g/ml, which can then be further diluted to other aqueous solutions. |
| Biological Activity | The activity of sVEGFR-1D1-4 was determined by its ability to abolish the binding of iodinated VEGF to solid surfaces or cell surfaces, and in Far-Western and cross-linking experiments with iodinated VEGF. |
| Storage | Lyophilized FLT-1 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution FLT1 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles. |

GENE INFORMATION

| | |
|------------------|--|
| Gene Name | FLT1 fms-related tyrosine kinase 1 (vascular endothelial growth factor/vascular permeability factor receptor) [Homo sapiens] |
| Synonyms | FLT1; fms-related tyrosine kinase 1 (vascular endothelial growth factor/vascular permeability factor receptor); FLT; VEGFR1; vascular endothelial growth factor receptor 1; EC 2.7.10.1; VEGFR-1; Vascular permeability factor receptor; Flt-1; Tyrosine-protein kinase receptor FLT; Tyrosine-protein kinase FRT; Fms-like tyrosine |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


| | |
|----------------------------|--|
| | kinase 1; FRT |
| Gene ID | 2321 |
| mRNA Refseq | NM_001159920 |
| Protein Refseq | NP_001153392 |
| MIM | 165070 |
| UniProt ID | P17948 |
| Chromosome Location | 13q12 |
| Pathway | Cytokine-cytokine receptor interaction; Endocytosis; Focal adhesion; Signaling by VEGF |
| Function | ATP binding; growth factor binding; identical protein binding; nucleotide binding; receptor activity; transferase activity; vascular endothelial growth factor receptor activity |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

PDB rendering based
on 1ft.



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA