

Recombinant Human FURIN, His-tagged

Cat. No. FURIN-13036H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human FURIN protein, fused to His-tag, was expressed in E.coli and purified by Ni-sepharose.
Species	Human
Source	E.coli
ProteinLength	C-term-300a.a.
Description	<p>The protein encoded by this gene belongs to the subtilisin-like proprotein convertase family. The members of this family are proprotein convertases that process latent precursor proteins into their biologically active products. This encoded protein is a calcium-dependent serine endoprotease that can efficiently cleave precursor proteins at their paired basic amino acid processing sites. Some of its substrates are: parathyroid hormone, transforming growth factor beta 1 precursor, proalbumin, pro-beta-secretase, membrane type-1 matrix metalloproteinase, beta subunit of pro-nerve growth factor and von Willebrand factor. It is also thought to be one of the proteases responsible for the activation of HIV envelope glycoproteins gp160 and gp140. This gene is thought to play a role in tumor progression. The use of alternate polyadenylation sites has been found for this gene.</p>
Storage	The protein is stored in PBS buffer at -20°C. Avoid repeated freezing and thawing cycles.
Storage Buffer	1M PBS (58mM Na ₂ HPO ₄ , 17mM NaH ₂ PO ₄ , 68mM NaCl, pH8.) added with 300mM

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Imidazole and 0.7% Sarcosyl, 15%glycerol.

GENE INFORMATION

Gene Name	FURIN furin (paired basic amino acid cleaving enzyme) [Homo sapiens]
Official Symbol	FURIN
Synonyms	FURIN; furin (paired basic amino acid cleaving enzyme); FUR, PACE, paired basic amino acid cleaving enzyme (furin, membrane associated receptor protein) , PCSK3; furin; SPC1; FES upstream region; dibasic processing enzyme; dibasic-processing enzyme; furin, membrane associated receptor protein; proprotein convertase subtilisin/kexin type 3; paired basic amino acid residue-cleaving enzyme; FUR; PACE; PCSK3;
Gene ID	5045
mRNA Refseq	NM_002569
Protein Refseq	NP_002560
MIM	136950
UniProt ID	P09958
Chromosome Location	15q26.1
Pathway	Activation of Matrix Metalloproteinases, organism-specific biosystem; Degradation of the extracellular matrix, organism-specific biosystem; Delta-Notch Signaling Pathway, organism-specific biosystem; Developmental Biology, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Extracellular matrix organization, organism-specific biosystem; Gamma-carboxylation, transport, and amino-terminal cleavage of proteins, organism-specific biosystem; Glypican 3 network, organism-specific biosystem;

Function

endopeptidase activity; metal ion binding; nerve growth factor binding; peptidase activity; peptide binding; protease binding; serine-type endopeptidase activity; serine-type endopeptidase activity; serine-type endopeptidase inhibitor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA