

Active Recombinant Mouse Fc Receptor, LgG, Low Affinity III, His-tagged

Cat. No. Fcgr3-488M **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Recombinant Mouse Fcgr3 is expressed in Murine myeloma cell line, which has a molecule weight of 22.5kDa. Ala31 Thr215, with a Cterminal 10His tag.

Species

Mouse

Source

Mammalian Cells

Description

Receptors for the Fc region of IgG (Fc γ Rs) are members of the Ig superfamily that function in the activation or inhibition of immune responses such as degranulation, phagocytosis, ADCC (antibody dependent cellular toxicity), cytokine release, and B cell proliferation. The Fc γ Rs have been divided into three classes based on close relationships in their extracellular domains; these groups are designated Fc γ RI (also known as CD64), The coordinated functioning of activating and inhibitory receptors is necessary for successful initiation, amplification, and termination of immune responses. Mouse CD16 is encoded by a single gene. The protein product is a type I transmembrane protein having two extracellular Ig-like domains. It is expressed on a variety of myeloid and lymphoid cells and associates with Fc γ R2 to deliver an activating signal upon ligand binding. Mouse CD32 is closely related to mouse CD16 throughout its extracellular domain (95% amino acid sequence identity), but has a divergent cytoplasmic domain and functions as an inhibitory receptor. Together these proteins constitute an activating/inhibiting receptor pair to regulate immune responses.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Form	Supplied as a 0.2 µmfiltered solution in PBS.
N-terminal Sequence	Ala31
Molecular Weight	22.5 kDa
Activity	Measured by its abilityto bind mouse IgG with an estimated KD < 15 nM.
Purity	>95%, by SDSPAGEunder reducing conditions and visualized by silver stain.
Endotoxin Level	<1.0 EU per 1 µg ofthe protein by the LAL method.
Storage	Use a manual defrostfreezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt,-20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution.3 months, -20 to -70 °C under sterile conditions after reconstitution.
OfficialSymbol	Fcgr3

GENE INFORMATION

Gene Name	Fcgr3 Fc receptor, IgG, low affinityIII [Mus musculus]
Synonyms	Fcgr3; Fc receptor, IgG,low affinity III; CD16; fcRIII; low affinity immunoglobulin gamma Fc region receptorIII; FcgammaRIII; fc-gamma RIII; Fcg receptor III; igG Fc receptor III; Fc gammareceptor III
Gene ID	14131
mRNA Refseq	NM_010188

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq	NP_034318
Chromosome Location	1 H3; 1 92.3 cM
Pathway	Adaptive Immune System; Immune System; Immunoregulatory interactions between a Lymphoid and a non-Lymphoid cell; Leishmaniasis; Natural killer cell mediated cytotoxicity; Osteoclast differentiation; Phagosome; Staphylococcus aureus infection; Systemic lupus erythematosus; Tuberculosis
Function	IgG binding; IgG receptor activity; receptor activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA