

## Recombinant Rat Fgf18 protein

**Cat. No.** Fgf18-588R    **Lot. No.** (See product label)

### SPECIFICATION

<b>Product Overview</b>	Recombinant Rat Fgf18 protein was expressed in Escherichia coli.
<b>Species</b>	Rat
<b>Source</b>	E.coli
<b>ProteinLength</b>	180
<b>Description</b>	<p>Rat FGF-18 is encoded by the FGF18 gene in rats. By phylogenetic analysis and gene location analysis, FGF-18 is divided into FGF-8 subfamily which has three members FGF-8, FGF-17 and FGF-18. Using FGF knockout mice model, the numbers of this subfamily were testified that have crucial roles of in embryo development. FGF-18<sup>-/-</sup> mice have decreased expression of osteogenic markers and delayed long-bone ossification. FGF-18 also has significant roles in lung development and has an anabolic effect on cartilage formation. Additionally, it has been shown in vitro that this protein is able to induce neurite outgrowth in PC12 cells. Rat FGF-18 shares 98 % amino acid sequence identity with human.</p>
<b>Form</b>	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4, 500 mM NaCl.
<b>Bio-activity</b>	Fully biologically active when compared to standard. The ED50 as determined by thymidine uptake assay using FGF-receptors transfected BaF3 cells is less than 0.5 ng/ml, corresponding to a specific activity of > 2.0 × 10 <sup>6</sup> IU/mg.

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Molecular Mass</b>	Approximately 21.0 kDa, a single non-glycosylated polypeptide chain containing 180 amino acids.
<b>AA Sequence</b>	EENVDFRIHVENQTRARDDVSRKQLRLYLQYSRTSGKHIQVLGRRISARGEDGDKY AQLLVETDTFGSQVRIKGGKETFYLCMNRKGLVGPDPGTSKECVFIEKVLNNYTA LMSAKYSGWYVGFTHKGRPRKGPKTRENQQDVHFMKRYPKGQTELQKPFKYTTVT KRSRRIRPHTPG
<b>Endotoxin</b>	Less than 1 EU/g of rRtFGF-18 as determined by LAL method.
<b>Purity</b>	>95% by SDS-PAGE and HPLC analysis.
<b>Storage</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
<b>Reconstitution</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

## GENE INFORMATION

<b>Gene Name</b>	<b>Fgf18</b>
<b>Official Symbol</b>	<b>Fgf18</b>
<b>Synonyms</b>	FGF18; fibroblast growth factor 18; FGF-18;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


Gene ID 29369

mRNA Refseq NM\_019199

Protein Refseq NP\_062072

UniProt ID O88182

SDS-PAGE of Fgf18-588R



 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA