

# Recombinant Mouse Fgfr4 Protein, His-tagged, FITC conjugated

**Cat. No.** Fgfr4-709MF    **Lot. No.** (See product label)

## SPECIFICATION

### Product Overview

FITC conjugated recombinant Mouse Fgfr4 (Leu17-Asp366) protein was fused to His-tag at C-terminus and expressed in human 293 cells (HEK293).

### Species

Mouse

### Source

HEK293

### ProteinLength

Leu17-Asp366

### Description

Fibroblast growth factor receptor 4(FGFR4) is also known as CD334, JTK2, hydroxyaryl-protein kinase, TKF, protein-tyrosine kinase . The FGFR4 gene provides instructions for making a protein called fibroblast growth factor receptor 4. This protein is part of a family of fibroblast growth factor receptors that share similar structures and functions. These receptor proteins play a role in important processes such as cell division, regulating cell growth and maturation, formation of blood vessels, wound healing, and embryo development.The FGFR4 protein interacts with specific growth factors to conduct signals from the environment outside the cell to the nucleus. The nucleus responds to these signals by switching on or off appropriate genes that help the cell adjust to changes in the environment. In response, the cell might divide, move, or mature to take on specialized functions. Although specific functions of FGFR4 remain unclear, studies indicate that the gene is involved in muscle development and the maturation of bone cells in the skull. The FGFR4 gene may also play a role in the development and maintenance of specialized cells (called

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	foveal cones) in the light-sensitive layer (the retina) at the back of the eye.
<b>Form</b>	Lyophilized
<b>Molecular Mass</b>	The protein has a calculated MW of 40.7 kDa. The protein migrates as 55-67 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.
<b>N-terminal Sequence Analysis</b>	Leu 17
<b>Endotoxin</b>	< 1.0 EU/ µg by the LAL method.
<b>Purity</b>	> 90 % as determined by SDS-PAGE
<b>Characteristic</b>	Disulfide-linked homodimer Labeled with FITC via amines Excitation source: 488 nm spectral line, argon-ion laser Excitation Wavelength: 488 nm Emission Wavelength: 535 nm
<b>Storage</b>	For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles. This product is stable after storage at: -20 to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution.
<b>Storage Buffer</b>	Lyophilized from 0.22 µm filtered solution in PBS, pH7.4, 10% trehalose.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 µg/µL. Centrifuge the vial at 4 centigrade before opening to recover the entire

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

contents.

**Conjugation** FITC**GENE INFORMATION****Gene Name** Fgfr4**Official Symbol** Fgfr4**Synonyms** FGFR4; fibroblast growth factor receptor 4; CTLA-2-beta protein; protein-tyrosine kinase receptor MPK-11; fibroblast growth factor receptor 4 16 minus form; Fgfr-4**Gene ID** 14186**mRNA Refseq** NM\_008011**Protein Refseq** NP\_032037**UniProt ID** Q03142 Tel: 1-631-559-9269 1-516-512-3133 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA