

Recombinant Human C2orf3 293 Cell Lysate

Cat. No. GCFC2-8082HCL Lot. No. (See product label)

SPECIFICATION

| | |
|----------------------------|---|
| Species | Human |
| Source | HEK293 |
| Description | Antigen standard for chromosome 2 open reading frame 3 (C2orf3) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection. |
| Components | This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol). |
| Size | 0.1 mg |
| Storage Instruction | Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment. |
| Applications | ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

| | |
|----------------------------|---|
| Gene Name | GCFC2 GC-rich sequence DNA-binding factor 2 [Homo sapiens] |
| Official Symbol | GCFC2 |
| Synonyms | GCF; TCF9; DNABF; C2orf3; GC-rich sequence DNA-binding factor 2; TCF-9; GC binding factor; GC binding factor; transcription factor 9 (binds GC-rich sequences) |
| Gene ID | 6936 |
| mRNA Refseq | NM_001201334 |
| Protein Refseq | NP_001188263 |
| MIM | 189901 |
| UniProt ID | P16383 |
| Chromosome Location | 2p12 |
| Function | RNA polymerase II core promoter proximal region sequence-specific DNA binding; RNA polymerase II core promoter proximal region sequence-specific DNA binding transcription factor activity involved in negative regulation of transcription |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA