

Recombinant Human GPX4 Protein, His-tagged

Cat. No. GPX4-027H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active recombinant N-terminal His-tagged full length human GPX4 protein (1-170) was expressed in E. coli strain C321.ΔA.exp.
Species	Human
Source	E.coli
ProteinLength	1-170
Description	<p>The protein encoded by this gene belongs to the glutathione peroxidase family, members of which catalyze the reduction of hydrogen peroxide, organic hydroperoxides and lipid hydroperoxides, and thereby protect cells against oxidative damage. Several isozymes of this gene family exist in vertebrates, which vary in cellular location and substrate specificity. This isozyme has a high preference for lipid hydroperoxides and protects cells against membrane lipid peroxidation and cell death. It is also required for normal sperm development; thus, it has been identified as a 'moonlighting' protein because of its ability to serve dual functions as a peroxidase, as well as a structural protein in mature spermatozoa. Mutations in this gene are associated with Sedaghatian type of spondylometaphyseal dysplasia (SMDS). This isozyme is also a selenoprotein, containing the rare amino acid selenocysteine (Sec) at its active site. Sec is encoded by the UGA codon, which normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, designated the Sec insertion sequence (SECIS) element, that is necessary for the recognition of UGA as a Sec codon, rather</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

than as a stop signal. Transcript variants resulting from alternative splicing or use of alternate promoters have been described to encode isoforms with different subcellular localization.

Molecular Mass 21.2 kDa

Purity ≥65% estimated by SDS-PAGE

Unit Definition One unit is defined as the amount of enzyme required to produce 1 μmol of NADP+ per minute at 25 centigrade in 50 mM Tris-HCl, pH 7.6, with 5 mM EDTA, 1 mM GSH, 0.076 units glutathione reductase, 263 μM NADPH and 0.5 mM of cumene hydroperoxide.

Applications WB, ELISA, and enzymatic assays

Stability ≥ 9 months

Storage At -80 centigrade.

Storage Buffer 50 mM potassium phosphate buffer, pH 7.6, with 0.1 mM DTT and 5% glycerol

GENE INFORMATION

Gene Name GPX4 glutathione peroxidase 4 [Homo sapiens (human)]

Official Symbol GPX4


Synonyms GPX4; glutathione peroxidase 4; MCSP; SMDS; GPx-4; PHGPx; snGPx; GSHPx-4; snPHGPx; phospholipid hydroperoxide glutathione peroxidase; epididymis secretory sperm binding protein; phospholipid hydroperoxidase; phospholipid hydroperoxide glutathione peroxidase, mitochondrial; selenoprotein GPX4; sperm nucleus

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

glutathione peroxidase; EC 1.11.1.12

Gene ID 2879**mRNA Refseq** NM_002085**Protein Refseq** NP_002076**MIM** 138322**UniProt ID** P36969**SDS-PAGE analysis
of GPX4**


Lane 1: MW Markers

Lane 2: GPX4 (2 µg)

Lane 3: GPX4 (4 µg)

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA

GPX4 Activity



GPX4 activity was determined using Glutathione Peroxidase Assay Kit.

GPX4 protein has a selenocysteine incorporated in the active site that has been confirmed by mass spectrometry. Selenocysteine is incorporated in the active site the UGA stop codon indicated in teal.

10	20	30	40	50
GSSHHHHHS	QDPNSMCASR	DDWRCARSMH	EFSAKDIDGH	MVNLDKYRGF
60	70	80	90	100
VCIVTNVASQ	UGKTEVNYTQ	LVDLHARYAE	CGLRILAFPC	NQFGKQEPGS
110	120	130	140	150
NEEIKEFAAG	YNVKFDMFSK	ICVNGDDAHP	LWKWMKIQPK	GKGILGNAIK
160	170	180		
WNFTKFLIDK	NGCVVKRYGP	MEEPLVIEKD	LPHYF	

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA