

Recombinant Human GRM2 293 Cell Lysate

Cat. No. GRM2-5736HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for glutamate receptor, metabotropic 2 (GRM2), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	GRM2 glutamate receptor, metabotropic 2 [Homo sapiens]
Official Symbol	GRM2
Synonyms	GRM2; glutamate receptor, metabotropic 2; metabotropic glutamate receptor 2; GPRC1B; mGlu2; MGLUR2; glutamate receptor homolog; glutamate metabotropic receptor 2; GLUR2;
Gene ID	2912
mRNA Refseq	NM_000839
Protein Refseq	NP_000830
MIM	604099
UniProt ID	Q14416
Chromosome Location	3
Pathway	Class C/3 (Metabotropic glutamate/pheromone receptors), organism-specific biosystem; Cocaine addiction, organism-specific biosystem; Cocaine addiction, conserved biosystem; GPCR ligand binding, organism-specific biosystem; GPCRs, Class C Metabotropic glutamate, pheromone, organism-specific biosystem; Glutamatergic synapse, organism-specific biosystem; Glutamatergic synapse, conserved biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

G-protein coupled receptor activity; calcium channel regulator activity; glutamate receptor activity; group II metabotropic glutamate receptor activity; receptor activity; signal transducer activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA