

## Recombinant Human GSTO2 293 Cell Lysate

Cat. No. GSTO2-5710HCL Lot. No. (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	Antigen standard for glutathione S-transferase omega 2 (GSTO2) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
<b>Applications</b>	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

<b>Gene Name</b>	GSTO2 glutathione S-transferase omega 2 [ Homo sapiens ]
<b>Official Symbol</b>	GSTO2
<b>Synonyms</b>	GSTO2; glutathione S-transferase omega 2; glutathione S-transferase omega-2; GSTO-2; MMA(V) reductase; monomethylarsonic acid reductase; glutathione S-transferase omega 2-2; glutathione-S-transferase-like protein; bA127L20.1 (novel glutathione-S-transferase); glutathione-dependent dehydroascorbate reductase; GSTO 2-2; bA127L20.1;
<b>Gene ID</b>	119391
<b>mRNA Refseq</b>	NM_001191013
<b>Protein Refseq</b>	NP_001177942
<b>MIM</b>	612314
<b>UniProt ID</b>	Q9H4Y5
<b>Chromosome Location</b>	10q25.1
<b>Pathway</b>	Biological oxidations, organism-specific biosystem; Drug metabolism - cytochrome P450, organism-specific biosystem; Drug metabolism - cytochrome P450, conserved biosystem; Glutathione conjugation, organism-specific biosystem; Glutathione metabolism, organism-specific biosystem; Glutathione metabolism, conserved

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



biosystem; Metabolism, organism-specific biosystem;

**Function**

glutathione dehydrogenase (ascorbate) activity; glutathione transferase activity;  
glutathione transferase activity; methylarsonate reductase activity; oxidoreductase  
activity; transferase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA