

Active Recombinant Mouse Gdf15 protein

Cat. No. Gdf15-393M Lot. No. (See product label)

SPECIFICATION

Product Overview Recombinant Mouse Gdf15(Ser189-Ala303) was expressed in E. coli.

Species Mouse

Source E.coli

ProteinLength Ser189-Ala303

Description

Growth Differentiation Factor 15 (GDF-15), also called Macrophage Inhibitory Cytokine 1 (MIC-1), Placental Transforming Growth Factor beta, Prostate-derived Factor, and Placental Bone Morphogenetic Protein, is a divergent member of the TGF-beta superfamily. Cellular responses to TGF-beta proteins are mediated by hetero-oligomeric complexes of type I and type II serine/threonine kinase receptors. GDF-15 is highly expressed in placenta and brain, and it is expressed at lower levels in kidney, pancreas, prostate, and colon. Similar to other TGF-beta family proteins, GDF-15 is synthesized as a large precursor protein that is cleaved at a dibasic cleavage site (RxxR) to release the mature protein. Mature mouse GDF-15 shares 66% and 97% amino acid sequence identity with the human and rat proteins, respectively. The C-terminal domain of GDF-15 contains seven characteristic conserved cysteine residues necessary for the formation of the cysteine knot and the single inter-chain disulfide bond. Biologically active GDF-15 is a disulfide-linked homodimer of the mature protein. GDF-15 has been shown to have various functions, including inhibition of TNF-alpha production from lipopolysaccharide-stimulated macrophages and the induction of cartilage formation. GDF-15 also promotes

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

neuronal survival, and hypothalamic expression of GDF-15 causes appetite suppression via modulation of neuropeptide Y and pro-opiomelanocortin levels. GDF-15 is cardioprotective via inhibition of platelet activation, limiting atherosclerosis, promoting recovery following myocardial infarction, and regulating angiogenesis. Exposure of cardiomyocytes to GDF-15 results in Smad2 and Smad3 phosphorylation.

Predicted N Terminal	Ser189
Form	Lyophilized from a 0.2 µm filtered solution in HCl.
Bio-activity	Measured by its binding ability in a functional ELISA. When Recombinant Mouse GSF-15 is immobilized at 2 g/mL (100 µ/well), the concentration of Recombinant Mouse Activin RIB/ALK-4 Fc Chimera that produces 50% of the optimal binding response is approximately 1-6 µg/mL.
Molecular Mass	Predicted Molecular Mass: 13 kDa; SDS-PAGE: 9-13 kDa, reducing conditions.
Endotoxin	<0.10 EU per 1 µg of the protein by the LAL method.
Purity	>90%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	Reconstitute at 250 µg/mL in 4 mM HCl.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

GENE INFORMATION

Gene Name Gdf15 growth differentiation factor 15 [Mus musculus]

Official Symbol Gdf15

Synonyms GDF15; growth differentiation factor 15; growth/differentiation factor 15; GDF-15; macrophage inhibiting cytokine-1; SBF; MIC-1; NAG-1;

Gene ID 23886

mRNA Refseq NM_011819

Protein Refseq NP_035949

UniProt ID Q9Z0J7

Chromosome Location 8; 8 B3.3

Function growth factor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA