

Recombinant Human HAO1 293 Cell Lysate

Cat. No. HAO1-5638HCL Lot. No. (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for hydroxyacid oxidase (glycolate oxidase) 1 (HAO1) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	HAO1 hydroxyacid oxidase (glycolate oxidase) 1 [Homo sapiens]
Official Symbol	HAO1
Synonyms	HAO1; hydroxyacid oxidase (glycolate oxidase) 1; GOX1; hydroxyacid oxidase 1; GOX; glycolate oxidase; (S)-2-hydroxy-acid oxidase; HAOX1; MGC142225; MGC142227;
Gene ID	54363
mRNA Refseq	NM_017545
Protein Refseq	NP_060015
MIM	605023
UniProt ID	Q9UJM8
Chromosome Location	20p12
Pathway	Glyoxylate and dicarboxylate metabolism, organism-specific biosystem; Glyoxylate and dicarboxylate metabolism, conserved biosystem; Glyoxylate metabolism, organism-specific biosystem; Metabolic pathways, organism-specific biosystem; Metabolism, organism-specific biosystem; Metabolism of amino acids and derivatives, organism-specific biosystem; Peroxisome, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Function

(S)-2-hydroxy-acid oxidase activity; FMN binding; glycolate oxidase activity; glycolate oxidase activity; glyoxylate oxidase activity; long-chain-(S)-2-hydroxy-long-chain-acid oxidase activity; medium-chain-(S)-2-hydroxy-acid oxidase activity; oxidoreductase activity; very-long-chain-(S)-2-hydroxy-acid oxidase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA