

Recombinant Human HDAC4 cell lysate

Cat. No. HDAC4-001HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human HDAC4 (aa 612-1084) derived in Baculovirus-Insect cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Insect Cells
ProteinLength	612-1084 a.a.
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[HDAC4 histone deacetylase 4 \[Homo sapiens \]](#)

Official Symbol

HDAC4

Synonyms

HDAC4; histone deacetylase 4; HA6116; HD4; HDAC 4; HDAC A; HDACA; KIAA0288; histone deacetylase A; AHO3; BDMR; HDAC-4; HDAC-A;

Gene ID

[9759](#)

mRNA Refseq

[NM_006037](#)

Protein Refseq

[NP_006028](#)

MIM

[605314](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	P56524
Chromosome Location	2q37.3
Pathway	Cell cycle, organism-specific biosystem; Endochondral Ossification, organism-specific biosystem; MicroRNAs in cardiomyocyte hypertrophy, organism-specific biosystem; NOTCH1 Intracellular Domain Regulates Transcription, organism-specific biosystem; Signal Transduction, organism-specific biosystem; Signaling by NOTCH, organism-specific biosystem; Signaling by NOTCH1, organism-specific biosystem;
Function	NAD-dependent histone deacetylase activity (H3-K14 specific); NAD-dependent histone deacetylase activity (H3-K9 specific); NAD-dependent histone deacetylase activity (H4-K16 specific); activating transcription factor binding; histone deacetylase activity; histone deacetylase activity (H3-K16 specific); histone deacetylase binding; hydrolase activity; potassium ion binding; protein binding; protein deacetylase activity; protein kinase binding; repressing transcription factor binding; contributes_to sequence-specific DNA binding; transcription factor binding; contributes_to transcription regulatory region DNA binding; zinc ion binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA