

## Recombinant Human HDAC6 293 Cell Lysate

Cat. No. HDAC6-5603HCL Lot. No. (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	Antigen standard for histone deacetylase 6 (HDAC6) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
<b>Applications</b>	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

**Gene Name** HDAC6 histone deacetylase 6 [ Homo sapiens ]

**Official Symbol** HDAC6

**Synonyms** HDAC6; histone deacetylase 6; FLJ16239; HD6; JM21; KIAA0901;

**Gene ID** 10013

**mRNA Refseq** NM\_006044

**Protein Refseq** NP\_006035

**MIM** 300272

**UniProt ID** Q9UBN7

**Chromosome Location** Xp11.23

**Pathway** Cell cycle, organism-specific biosystem; NOTCH1 Intracellular Domain Regulates Transcription, organism-specific biosystem; Signal Transduction, organism-specific biosystem; Signaling by NOTCH, organism-specific biosystem; Signaling by NOTCH1, organism-specific biosystem; Signaling events mediated by HDAC Class I, organism-specific biosystem; Signaling events mediated by HDAC Class II, organism-specific biosystem;

**Function** Hsp90 protein binding; NAD-dependent histone deacetylase activity (H3-K14

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



specific); NAD-dependent histone deacetylase activity (H3-K9 specific); NAD-dependent histone deacetylase activity (H4-K16 specific); actin binding; alpha-tubulin binding; beta-catenin binding; beta-tubulin binding; dynein complex binding; enzyme binding; histone deacetylase activity; histone deacetylase activity (H3-K16 specific); histone deacetylase binding; hydrolase activity; metal ion binding; microtubule binding; microtubule binding; polyubiquitin binding; protein binding; tau protein binding; tubulin deacetylase activity; tubulin deacetylase activity; zinc ion binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA