

## Recombinant Human HDAC9 293 Cell Lysate

**Cat. No.** HDAC9-5601HCL    **Lot. No.** (See product label)

### SPECIFICATION

<b>Species</b>	Human
<b>Source</b>	HEK293
<b>Description</b>	Antigen standard for histone deacetylase 9 (HDAC9), transcript variant 3 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
<b>Components</b>	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
<b>Size</b>	0.1 mg
<b>Storage Instruction</b>	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
<b>Applications</b>	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

## GENE INFORMATION

**Gene Name** [HDAC9 histone deacetylase 9 \[ Homo sapiens \]](#)

**Official Symbol** HDAC9

**Synonyms** HDAC9; histone deacetylase 9; HD7; HDAC; HDAC7B; KIAA0744; MITR; histone deacetylase 7B; histone deacetylase 4/5-related protein; MEF-2 interacting transcription repressor (MITR) protein; HD9; HD7b; HDRP; HDAC7; HDAC9B; HDAC9FL; DKFZp779K1053;

**Gene ID** [9734](#)

**mRNA Refseq** [NM\\_001204144](#)

**Protein Refseq** [NP\\_001191073](#)

**MIM** [606543](#)

**UniProt ID** [Q9UKV0](#)

**Chromosome Location** 7p21.1

**Pathway** MicroRNAs in cardiomyocyte hypertrophy, organism-specific biosystem; NOTCH1 Intracellular Domain Regulates Transcription, organism-specific biosystem; Signal Transduction, organism-specific biosystem; Signaling by NOTCH, organism-specific biosystem; Signaling by NOTCH1, organism-specific biosystem; Signaling events mediated by HDAC Class I, organism-specific biosystem; Signaling events mediated

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

by HDAC Class II, organism-specific biosystem;

**Function**

NAD-dependent histone deacetylase activity (H3-K14 specific); NAD-dependent histone deacetylase activity (H3-K9 specific); NAD-dependent histone deacetylase activity (H4-K16 specific); histone deacetylase activity; NOT histone deacetylase activity; histone deacetylase activity (H3-K16 specific); histone deacetylase binding; histone deacetylase binding; hydrolase activity; protein binding; protein deacetylase activity; protein kinase C binding; repressing transcription factor binding; repressing transcription factor binding; repressing transcription factor binding; transcription corepressor activity; transcription factor binding; transcription factor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA