

Recombinant Human HDGF 293 Cell Lysate

Cat. No. HDGF-5598HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for hepatoma-derived growth factor (high-mobility group protein 1-like) (HDGF), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	HDGF hepatoma-derived growth factor [Homo sapiens]
Official Symbol	HDGF
Synonyms	HDGF; hepatoma-derived growth factor; hepatoma derived growth factor (high mobility group protein 1 like); high mobility group protein 1 like; HMG1L2; HMG-1L2; high mobility group protein 1-like 2; FLJ96580; DKFZp686J1764;
Gene ID	3068
mRNA Refseq	NM_001126050
Protein Refseq	NP_001119522
MIM	600339
UniProt ID	P51858
Chromosome Location	1q21-q23
Pathway	Activation of Chaperone Genes by XBP1(S), organism-specific biosystem; Activation of Chaperones by IRE1alpha, organism-specific biosystem; Diabetes pathways, organism-specific biosystem; Disease, organism-specific biosystem; Unfolded Protein Response, organism-specific biosystem;
Function	DNA binding; growth factor activity; heparin binding; nucleotide binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA