

Recombinant Human HIST2H2AC Protein, His-tagged

Cat. No. HIST2H2AC-2241H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human HIST2H2AC Protein (Met1-Lys129) with a N-His tag was expressed in E. coli.
Species	Human
Source	E.coli
ProteinLength	Met1-Lys129
Description	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H2A family.
Form	Freeze-dried powder
Molecular Mass	Predicted Molecular Mass: 17.7 kDa Accurate Molecular Mass: 18 kDa
Endotoxin	<1.0 EU per 1g (determined by the LAL method).
Purity	> 90%

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications	Positive Control; Immunogen; SDS-PAGE; WB.
Stability	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 centigrade for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Storage	Avoid repeated freeze/thaw cycles. Store at 2-8 centigrade for one month. Aliquot and store at -80 centigrade for 12 months.
Storage Buffer	20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
Reconstitution	Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	HIST2H2AC histone cluster 2, H2ac [Homo sapiens (human)]
Official Symbol	HIST2H2AC
Synonyms	H2AC20; H2A clustered histone 20; H2A; H2A/q; H2AFQ; H2A-GL101; HIST2H2AC; histone H2A type 2-C; H2A histone family, member Q; histone 2, H2ac; histone H2A-GL101; histone H2A/q; histone IIa; histone cluster 2 H2A family member c; histone cluster 2, H2ac
Gene ID	8338
mRNA Refseq	NM_003517

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq NP_003508

MIM 602797

UniProt ID Q16777



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA