

Recombinant Mouse HMGB1 cell lysate

Cat. No. HMGB1-1677MCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mouse HMGB1 / HMG1 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Mouse
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[Hmgb1 high mobility group box 1 \[Mus musculus \]](#)

Official Symbol

HMGB1

Synonyms

HMGB1; high mobility group box 1; high mobility group protein B1; high mobility group protein 1; DEF; p30; Hmg1; HMG-1; SBP-1; amphoterin; MGC103168; MGC103169; MGC117896; MGC117897;

Gene ID

[15289](#)

mRNA Refseq

[NM_010439](#)

Protein Refseq

[NP_034569](#)

Pathway

Activated TLR4 signalling, organism-specific biosystem; Activation of DNA fragmentation factor, organism-specific biosystem; Androgen Receptor Signaling Pathway, organism-specific biosystem; Apoptosis, organism-specific biosystem; Apoptosis induced DNA fragmentation, organism-specific biosystem; Apoptotic

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

executionphase, organism-specific biosystem; Base excision repair, organism-specific biosystem;

Function

5S rRNA binding; DNA binding; DNA binding, bending; RAGE receptor binding; bent DNA binding; binding; calcium-dependent protein kinase regulator activity; crossed form four-way junction DNA binding; cytokine activity; cytokine activity; double-stranded DNA binding; four-way junction DNA binding; glycolipid binding; heparin binding; open form four-way junction DNA binding; peptide binding; phosphatidylserine binding; protein dimerization activity; protein kinase activator activity; repressing transcription factor binding; sequence-specific DNA binding transcription factor activity; single-stranded DNA binding; transcription factor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA