

Active Recombinant Human HMGB1 protein, His&hFc-tagged

Cat. No. HMGB1-454H Lot. No. (See product label)

SPECIFICATION

Product Overview	Active Recombinant Human HMGB1 protein(P09429)(1~215 aa), fused with N-terminal His tag and C-terminal hFc tag, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	1-215 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	<p>The binding activity of HMG1 with TP53.</p> <p>High Mobility Group Protein 1 (HMG1), also known as high mobility group box 1 protein belongs to high mobility group and contains HMG-box domain. HMG1 is one of the most important chromatin proteins. This nuclear protein organizes the DNA and regulates transcription. It supports transcription of many genes in interactions with many transcription factors. HMGB1 is secreted by immune cells (like macrophages, monocytes and dendritic cells) through leaderless secretory pathway. Activated macrophages and monocytes secrete HMGB1 as a cytokine mediator of Inflammation. Besides, Tumor Protein p53 (TP53) has been identified as an interactor of HMG1, thus a binding ELISA assay was conducted to detect the interaction of recombinant human HMG1 and recombinant human TP53. Briefly, HMG1 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100uL were</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

then transferred to TP53-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-HMG1 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µ stop solution to the wells and read at 450nm immediately. The binding activity of HMG1 and TP53 was shown, and this effect was in a dose dependent manner.

Molecular Mass 51.2kDa

Endotoxin <1.0EU per 1µg (determined by the LAL method).

Purity > 95% as determined by SDS-PAGE.

Storage Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.

Reconstitution Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name HMGB1 high mobility group box 1 [Homo sapiens]

Official Symbol HMGB1

Synonyms HMGB1; high mobility group box 1; high mobility group (nonhistone chromosomal) protein 1, high mobility group box 1, HMG1; high mobility group protein B1; Amphoterin; DKFZp686A04236; high mobility group protein 1; HMG3; SBP 1; Sulfoglucuronyl carbohydrate binding protein; HMG-1; high-mobility group box 1; high-mobility group (nonhistone chromosomal) protein 1; HMG1; SBP-1;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID 3146


mRNA Refseq NM_002128

Protein Refseq NP_002119


MIM 163905

UniProt ID P09429

SDS-PAGE



Activity



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA