

Recombinant Human HSD17B4 293 Cell Lysate

Cat. No. HSD17B4-5373HCL **Lot. No.** (See product label)

SPECIFICATION

Species	Human
Source	HEK293
Description	Antigen standard for hydroxysteroid (17-beta) dehydrogenase 4 (HSD17B4) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [HSD17B4 hydroxysteroid \(17-beta\) dehydrogenase 4 \[Homo sapiens \]](#)

Official Symbol HSD17B4

Synonyms

HSD17B4; hydroxysteroid (17-beta) dehydrogenase 4; peroxisomal multifunctional enzyme type 2; 3 alpha; 7 alpha; 12 alpha trihydroxy 5 beta cholest 24 enoyl CoA hydratase; 17 beta HSD IV; 17 beta hydroxysteroid dehydrogenase 4; 17beta estradiol dehydrogenase type IV; beta hydroxyacyl dehydrogenase; beta keto reductase; D 3 hydroxyacyl CoA dehydratase; D bifunctional protein; peroxisomal; DBP; MFE 2; peroxisomal multifunctional protein 2; SDR8C1; short chain dehydrogenase/reductase family 8C; member 1; 17-beta-HSD 4; 17-beta-HSD IV; beta-keto-reductase; multifunctional protein 2; beta-hydroxyacyl dehydrogenase; D-3-hydroxyacyl-CoA dehydratase; D-bifunctional protein, peroxisomal; 17-beta-hydroxysteroid dehydrogenase 4; 17beta-estradiol dehydrogenase type IV; short chain dehydrogenase/reductase family 8C, member 1; 3-alpha,7-alpha,12-alpha-trihydroxy-5-beta-cholest-24-enoyl-CoA hydratase; MFE-2; MPF-2;

Gene ID [3295](#)

mRNA Refseq [NM_000414](#)

Protein Refseq [NP_000405](#)

MIM [601860](#)

UniProt ID [P51659](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	5q2
Pathway	Beta-oxidation of pristanoyl-CoA, organism-specific biosystem; Beta-oxidation of very long chain fatty acids, organism-specific biosystem; Bile acid and bile salt metabolism, organism-specific biosystem; Bile acid biosynthesis, cholesterol =>cholate, organism-specific biosystem; Bile acid biosynthesis, cholesterol => cholate, conserved biosystem;
Function	3-hydroxyacyl-CoA dehydrogenase activity; 3alpha,7alpha,12alpha-trihydroxy-5beta-cholest-24-enoyl-CoA hydratase activity; estradiol 17-beta-dehydrogenase activity; isomerase activity; long-chain-enoyl-CoA hydratase activity; lyase activity; nucleotide binding; oxidoreductase activity; sterol binding; sterol transporter activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA