

# Active Recombinant Human ICAM1 protein, Fc/Avi-tagged, Biotinylated

Cat. No. ICAM1 Lot. No. (See product label)

## SPECIFICATION

<b>Product Overview</b>	Biotinylated Recombinant Human ICAM1(Gln28-Glu480) protein, fused to Fc/Avi tag at the C-terminus, was expressed in HEK293 cells .
<b>Species</b>	Human
<b>Source</b>	HEK293
<b>ProteinLength</b>	Gln28-Glu480
<b>Description</b>	<p>ICAM1 (Intercellular Adhesion Molecule-1), also known as CD54, is a transmembrane cell adhesion glycoprotein and a member of the immunoglobulin supergene family. The ICAM sub-family consists of five members, ICAM-1 through ICAM-5, and they vary in their tissue expression and number of Ig-like domains in the extracellular domain (ECD). Full-length ICAM-1 contains five Ig-like domains in the ECD, a single transmembrane domain and short intracellular domain. Alternative splicing of ICAM-1 results in at least six membrane-bound forms and one soluble form (1). The ECD of mature, full-length human ICAM-1 shares 78% and 54% amino acid sequence identity with rat and mouse ICAM-1, respectively. ICAM-1 plays an important role in both innate and adaptive immune responses and is upregulated in endothelial and epithelial cells at sites of inflammation (2, 3). ICAM-1 mediates vascular adhesion and paracellular migration of leukocytes expressing activated LFA-1 (CD11a/CD18) and Mac-1 (CD11b/CD18) 2, 3). It also binds several non-integrin ligands including CD43/Sialophorin, Fibrinogen, Hyaluronan, rhinoviruses, and Plasmodium falciparum-</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

infected erythrocytes (4-8). Soluble ICAM-1 promotes angiogenesis and serves as an indicator of vascular endothelial cell activation or damage (9). Elevated levels of soluble ICAM-1 are associated with cardiovascular disease, type 2 diabetes, organ transplant dysfunction, oxidant stress, increased abdominal fat mass, hypertension, liver disease, certain malignancies, and cerebral malaria (10-18). More recently, ICAM-1 was shown to be a novel regulator of group 2 innate lymphoid cells (ILC2s) which play a key role in allergic airway inflammation (19).

**Predicted N Terminal** Gln28

**Form** Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose.

**Bio-activity** Measured by its binding ability in a functional ELISA. When Human ICAM1 Antibody is immobilized at 0.5 µg/mL (100 µL/well), Biotinylated Recombinant Human ICAM1 Fc Chimera Avi-tag binds with an ED50 of 3-18 ng/mL. Measured by its ability to support the adhesion of HSB2 human peripheral blood acute lymphoblastic leukemia cells. The ED50 for this effect is 0.3-2.7 µg/mL.

**Molecular Mass** 95-125 kDa, under reducing conditions

**Endotoxin** <0.10 EU per 1 µg of the protein by the LAL method.

**Purity** >95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.

**Applications** Bioactivity

**Storage** Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 3 months, -20 to -70 °C under sterile conditions after

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reconstitution.

**Reconstitution** Reconstitute at 500 µg/mL in PBS.

**Conjugation** Biotin

## GENE INFORMATION

**Gene Name** [ICAM1 intercellular adhesion molecule 1 \[ Homo sapiens \]](#)

**Official Symbol** [ICAM1](#)

**Synonyms** ICAM1; intercellular adhesion molecule 1; BB2; CD54; human rhinovirus receptor; ICAM-1; cell surface glycoprotein P3.58; major group rhinovirus receptor; intercellular adhesion molecule 1 (CD54), human rhinovirus receptor; P3.58;

**Gene ID** [3383](#)

**mRNA Refseq** [NM\\_000201](#)

**Protein Refseq** [NP\\_000192](#)

**MIM** [147840](#)

**UniProt ID** [P05362](#)

**SDS-PAGE** 2 µg/lane Protein was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by Coomassie® Blue staining.

**Binding Activity**

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA