

# Recombinant Human ICOS Protein, Gly/Pro-tagged, Alexa Fluor 488 conjugated

Cat. No. ICOS-390HAF488 Lot. No. (See product label)

## SPECIFICATION

### Product Overview

Alexa Fluor 488 conjugated recombinant human ICOS (Met 1-Phe 141) was expressed in HEK293.

### Species

Human

### Source

HEK293

### ProteinLength

Met1-Phe141

### Description

Inducible costimulator (ICOS), also called AILIM (activation-inducible lymphocyte immunomediatory molecule) is a cell-surface receptor, and belongs to the CD28 family of immune costimulatory receptors consisting of CD28, CTLA-4 and PD-1. The interaction of B7-H2/ICOS plays a critical role in Th cell differentiation, T-B cell interactions which is essential for germinal center formation, and humoral immune responses, and as well as the production of cytokine IL-4. In addition, ICOS is more potent in the induction of IL-10 production, a cytokine important for suppressive function of T regulatory cells. The B7-1/B7-2-CD28/CTLA-4 and ICOS-B7RP-1 pathway provides key second signals that can regulate the activation, inhibition and fine-tuning of T-lymphocyte responses. ICOS stimulates both Th1 and Th2 cytokine production but may have a preferential role in Th2 cell development. Moreover, The B7-1/B7-2-CD28/CTLA-4 and ICOS-B7RP-1 pathway has been suggested of being involved in the development of airway inflammation and airway hyperresponsiveness.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Form</b>	Lyophilized
<b>N-terminal Sequence Analysis</b>	Glu 21
<b>Endotoxin</b>	< 1.0 EU/ µg of the protein as determined by the LAL method.
<b>Purity</b>	> 94 % as determined by SDS-PAGE
<b>Characteristic</b>	Disulfide-linked homodimer Labeled with Alexa Fluor 488 via amines Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
<b>Stability</b>	Samples are stable for up to 12 months from date of receipt at -70 centigrade.
<b>Storage</b>	Store it under sterile conditions at -20 centigrade to -80 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
<b>Storage Buffer</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5%-8% trehalose and mannitol are added as protectants before lyophilization.
<b>Shipping</b>	Shipped at ambient temperature. Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice unless customers require otherwise.
<b>Conjugation</b>	Alexa Fluor 488

## GENE INFORMATION

**Gene Name** [ICOS inducible T-cell co-stimulator \[ Homo sapiens \]](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Official Symbol</b>	ICOS
<b>Synonyms</b>	ICOS; inducible T-cell co-stimulator; inducible T-cell costimulator; activation inducible lymphocyte immunomediatory molecule; AILIM; CD278; inducible costimulator; activation-inducible lymphocyte immunomediatory molecule; CVID1; MGC39850;
<b>Gene ID</b>	29851
<b>mRNA Refseq</b>	NM_012092
<b>Protein Refseq</b>	NP_036224
<b>MIM</b>	604558
<b>UniProt ID</b>	Q9Y6W8

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA