

Recombinant Human Inducible T-cell Co-stimulator Ligand, Fc Chimera

Cat. No. ICOSLG-2143H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human B7H2 extracellular domain (C-terminus fused with human IgG1 Fc) produced in mammalian cells contains 457 amino acids with predicted MW 51.21D x 2 (dimer), and 12 cysteines (Fc has 6). It also has 6 N-link glycosylation sites with MW increase up to 30 KD.
Species	Human
Source	Mammalian Cells
Description	B7H2 is ligand for the T-cell specific cell surface receptor ICOS and acts as a costimulatory signal for T-cell proliferation and differentiation. It induces also B-cell proliferation and differentiation in plasma cells.
Purity	>90%.
Recommended Storage Temperature	-70°C.

GENE INFORMATION

Gene Name	ICOSLG inducible T-cell co-stimulator ligand [Homo sapiens]
Synonyms	inducible T-cell co-stimulator ligand; B7H2; GL50; B7-H2; B7RP1; CD275; ICOSL; LICOS; B7RP-1; ICOS-L; KIAA0653; ICOSLG; ICOS ligand; B7 homolog 2; B7-like

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

protein GI50; B7-related protein 1; transmembrane protein B7-H2 ICOS ligand; B7-like protein; CD275 antigen

Gene ID [23308](#)

mRNA Refseq [NM_015259](#)

Protein Refseq [NP_056074](#)

MIM [605717](#)

UniProt ID [O75144](#)

Chromosome Location 21q22.3

Pathway Cell adhesion molecules (CAMs); Intestinal immune network for IgA production; Signaling in Immune system

Function receptor binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA