

Recombinant Human IDH1 (R132H mutation) Protein, His-tagged

Cat. No. IDH1-030H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Active recombinant C-terminal His-tagged Human IDH1 (R132H mutation) protein (2-414) was expressed in E. coli.

Species

Human

Source


E.coli

ProteinLength

2-414

Description

Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the cytoplasm and peroxisomes. It contains the PTS-1 peroxisomal targeting signal sequence. The presence of this enzyme in peroxisomes suggests roles in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2-oxoglutarate, namely the alpha-hydroxylation of phytanic acid. The cytoplasmic enzyme serves a significant role in cytoplasmic NADPH production. Alternatively

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

spliced transcript variants encoding the same protein have been found for this gene.

Molecular Mass 47.9 kDa

Purity ≥95% as determined by SDS-PAGE

Unit Definition One unit is defined as the amount of enzyme required to convert 1 nmol of NADPH to NADP+, using 15 mM α-ketoglutarate as a substrate, per minute at room temperature in 25 mM Tris-HCl, pH 7.5, 150 mM sodium chloride, and 5 mM MgCl₂.

Stability ≥ 2 years

Storage At -80 centigrade.

Storage Buffer 50 mM Tris-HCl, pH 7.5, containing 200 mM sodium chloride, 5 mM β-mercaptoethanol, and 10% glycerol

GENE INFORMATION

Gene Name IDH1 isocitrate dehydrogenase (NADP(+)) 1 [Homo sapiens (human)]

Official Symbol IDH1

Synonyms IDH1; isocitrate dehydrogenase (NADP(+)) 1; IDH; IDP; IDCD; IDPC; PICD; HEL-216; HEL-S-26; isocitrate dehydrogenase [NADP] cytoplasmic; NADP(+)-specific ICDH; NADP-dependent isocitrate dehydrogenase, cytosolic; NADP-dependent isocitrate dehydrogenase, peroxisomal; epididymis luminal protein 216; epididymis secretory protein Li 26; epididymis secretory sperm binding protein; isocitrate dehydrogenase (NADP(+)) 1, cytosolic; isocitrate dehydrogenase 1 (NADP+), soluble; oxalosuccinate decarboxylase; EC 1.1.1.42

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID 3417


mRNA Refseq NM_005896

Protein Refseq NP_005887

MIM 147700

UniProt ID O75874

**SDS-PAGE analysis
of IDH1**




Lane 1: MW Markers

Lane 2: IDH1 R132H (5 µg)

Lane 3: IDH1 R132H (2.5 µg)

Lane 4: IDH1 R132H (1.25 µg)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA