

Recombinant Full Length Human IDH1 Protein, C-Flag-tagged

Cat. No. IDH1-1116HFL Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Full Length Human IDH1 Protein, fused to Flag-tag at C-terminus, was expressed in Mammalian cells.
Species	Human
Source	Mammalian Cells
Description	<p>Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the cytoplasm and peroxisomes. It contains the PTS-1 peroxisomal targeting signal sequence. The presence of this enzyme in peroxisomes suggests roles in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2-oxoglutarate, namely the alpha-hydroxylation of phytanic acid. The cytoplasmic enzyme serves a significant role in cytoplasmic NADPH production. Alternatively spliced transcript variants encoding the same protein have been found for this gene.</p>
Form	25 mM Tris HCl, pH 7.3, 100 mM glycine, 10% glycerol.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	46.5 kDa
AA Sequence	<p>MSKKISGGSVVEMQGDDEMTRIIWELIKEKLIFPYVELDLHSYDLGIENRDATNDQVTK DAAEAIKKHNVG VKCATITPDEKRVVEEFKLMWQKSPNGTIRNILGGTVFREAICKNI PRLVSGWVKPIIIGRHAYGDQYR ATDFVVPGPGKVEITYTPSDGTQKVTYLVHNFEE GGGVAMGMYNQDKSIEDFAHSSFQMALSKGWPLYLS TKNTILKKYDGRFKDIFQEI YDKQYKSQFEAQKIWYEHRLIDDMVAQAMKSEGGFIWACKNYDGDVQSDS VAQG YGSLGMMTSVLVCPDGKTVEAESAHGTVTRHYRMYQKQGQETSTNPIASIFAWTRGL AHRAKLDNNK ELAFFANALEEVSIETIEAGFMTKDLAACIKGLPNVQRSDYLNTFEFMDKLGENLKIKL AQAKLTRTRPLEQKLISEEDLAANDILDYKDDDDKV</p>
Purity	> 80% as determined by SDS-PAGE and Coomassie blue staining.
Stability	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
Storage	Store at -80 centigrade.
Concentration	>50 ug/mL as determined by microplate BCA method.
Preparation	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Protein Pathways	Citrate cycle (TCA cycle), Glutathione metabolism, Metabolic pathways
Full Length	Full L.

GENE INFORMATION


Gene Name [IDH1 isocitrate dehydrogenase \(NADP\(+\)\) 1 \[Homo sapiens \(human\) \]](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Official Symbol	IDH1
Synonyms	IDH; IDP; IDCD; IDPC; PICD; HEL-216; HEL-S-26
Gene ID	3417
mRNA Refseq	NM_005896.4
Protein Refseq	NP_005887.2
MIM	147700
UniProt ID	O75874



Coomassie blue staining of purified IDH1 protein.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA