

Active Recombinant Pig IFN γ protein, His-tagged

Cat. No. IFN γ -3366P **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Pig IFN γ protein(P17803)(21~166 aa), fused with N-terminal His tag, was expressed in HEK293.
Species	Pig
Source	HEK293
ProteinLength	21-166 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	<p>IFN-g is a dimerized soluble cytokine that is the only member of the type II class of interferons. The importance of IFNγ in the immune system stems in part from its ability to inhibit viral replication directly and most importantly from its immunostimulatory and immunomodulatory effects. IFN-gamma dimers bind to IFN-gamma RI (alpha subunits) which then interact with IFN-gamma RII (beta subunits) to form the functional receptor complex of two alpha and two beta subunits. Inclusion of IFN-gamma RII increases the binding affinity for ligand and the efficiency of signal transduction. A functional binding ELISA assay was conducted to detect the interaction of recombinant pig IFN-g and recombinant human IFNγR1. Briefly, IFN-g was diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100 μl were then transferred to IFNγR1-coated microtiter wells and incubated for 1h at 37°C. Wells were washed with PBST and incubated for 1h with anti-IFN-g pAb, then</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

aspirated and washed 3 times. After incubation with HRP labelled secondary antibody for 1h at 37°C, wells were aspirated and washed 5 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50 µ stop solution to the wells and read at 450/630 nm immediately. The binding activity of recombinant pig IFN-g and recombinant human IFNgR1 was shown, the EC50 for this effect is 0.38 ug/mL.

Molecular Mass	17.7kDa
Endotoxin	<1.0EU per 1µg (determined by the LAL method).
Purity	> 90% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	IFNG interferon-gamma [<i>Sus scrofa</i>]
Official Symbol	IFNg
Synonyms	IFNG; interferon-gamma; interferon gamma; IFN-gamma;
Gene ID	396991
mRNA Refseq	NM_213948

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


Protein Refseq

NP_999113

SDS-PAGE



Activity



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA