

Recombinant Active Pig IL10 Protein, His-tagged(C-ter)

Cat. No. IL10-136P Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Active Pig IL10 Protein with His tag (C-ter) was expressed in E. coli.
Species	Pig
Source	E.coli
Description	<p>The protein encoded by this gene is a cytokine produced primarily by monocytes and to a lesser extent by lymphocytes. This cytokine has pleiotropic effects in immunoregulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II Ags, and costimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. This cytokine can block NF-kappa B activity, and is involved in the regulation of the JAK-STAT signaling pathway. Knockout studies in mice suggested the function of this cytokine as an essential immunoregulator in the intestinal tract. Mutations in this gene are associated with an increased susceptibility to HIV-1 infection and rheumatoid arthritis.[provided by RefSeq, May 2011]</p>
Form	Powder
Bio-activity	Determined by its ability to induce proliferation in MC/9-2 cells. The ED50 for this effect is < 5 ng/mL.
AA Sequence	<p>MSIKSENSCIHFPTSLPHMLRELRAAFGPVKSFFQTKDQMGDLLLLTGSLLLEDFKGYL GCQALSEMIQFYLEDVMPKAESDGEDIKEHVNSLGEKLTLLRRLRRCHQFLPCENK SKAVEEVKSAFSKLGQERGVYKAMGEFDIFINYIEAYMTMKMRKN</p>

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	Endotoxin level is less than 0.01 EU/μg of the protein, as determined by the LAL test.
Purity	> 98% (by SDS-PAGE)
Applications	SDS-PAGE
Notes	For laboratory research only, not for drug, diagnostic or other use.
Storage	Lyophilized protein should be stored at -20°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Storage Buffer	PBS (pH 7.4)
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.

GENE INFORMATION

Gene Name	IL10 interleukin 10 [<i>Sus scrofa</i>]
Official Symbol	IL10
Synonyms	IL10; interleukin 10; interleukin-10; cytokine synthesis inhibitory factor; CSIF; IL-10;
Gene ID	397106
mRNA Refseq	NM_214041

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq

NP_999206

SDS-PAGE



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA