

Recombinant Human IL10RA

Cat. No. IL10RA-29631TH **Lot. No.** (See product label)

SPECIFICATION

| | |
|---------------------------|--|
| Product Overview | Recombinant fragment, corresponding to amino acids 1-232 of Human IL10RA fused to the Fc region of human IgG1, with N-linked and O-linked oligosaccharides, and expressed in modified human 293 cells. |
| Species | Human |
| Protein Length | 1-232 a.a. |
| Description | The protein encoded by this gene is a receptor for interleukin 10. This protein is structurally related to interferon receptors. It has been shown to mediate the immunosuppressive signal of interleukin 10, and thus inhibits the synthesis of proinflammatory cytokines. This receptor is reported to promote survival of progenitor myeloid cells through the insulin receptor substrate-2/PI 3-kinase/AKT pathway. Activation of this receptor leads to tyrosine phosphorylation of JAK1 and TYK2 kinases. Two transcript variants, one protein-coding and the other not protein-coding, have been found for this gene. |
| Tissue specificity | Spleen, thymus, and PBMC. Weak expression in pancreas, skeletal muscle, brain, heart, and kidney. Placenta, lung, and liver showed intermediate levels. Monocytes, B-cells, large granular lymphocytes, and T-cells express high levels. |
| Form | Lyophilised: It is recommended that 0.5 ml of sterile phosphate-buffered saline be added to the vial. Following reconstitution short-term storage at 4°C is recommended, and longer-term storage of aliquots at -18 to -20°C. Repeated freeze thawing is not |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

| | |
|---------------------------------|---|
| | reco |
| Purity | >95% by SDS-PAGE |
| Storage buffer | Preservative: None Constituents: 10% Trehalose, 1% Human serum albumin, PBS |
| Storage | Store at +4°C. |
| Sequences of amino acids | Theoretical Sequence:HGTELPSPPSVWFEEAFFHHILHWTPIPN QSESTCYEVALLR YGIESWNSISNCSQTLSDLTAVTL DLYHSNGYRARVRAVDGSRHSNWTVTNTRFS VDEVTLTVG SVNLEIHNGFILGKIQLPRPKMAPANDTYESIFSHFRE YEIAIRKVPGN FTFTHKKVKHENFSLTSGEVGEFCVQVK PSVASRSNKGMWSKEECISLTRQYFTG IPKVDKKVEPK SCDKHTCPCPAPELLGGPSVFLFPPKPKDTLMISRT PEVTCVVV DVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY NSTYRVVSVLTVLHQDWLNGKE YKCRVSNKALPAPIEK TISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYP S DIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTV DKSRWQQGNVFCFSVMHE ALHNHYTQKSLSLSPGK |
| Sequence Similarities | Belongs to the type II cytokine receptor family. |
| GENE INFORMATION | |
| Gene Name | IL10RA interleukin 10 receptor, alpha [Homo sapiens] |
| Official Symbol | IL10RA |
| Synonyms | IL10RA; interleukin 10 receptor, alpha; IL10R; interleukin-10 receptor subunit alpha; CD210a; CDW210A; HIL 10R; |
| Gene ID | 3587 |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

| | |
|----------------------------|---|
| mRNA Refseq | NM_001558 |
| Protein Refseq | NP_001549 |
| MIM | 146933 |
| Uniprot ID | Q13651 |
| Chromosome Location | 11q23 |
| Pathway | Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Jak-STAT signaling pathway, organism-specific biosystem; Jak-STAT signaling pathway, conserved biosystem; Toxoplasmosis, organism-specific biosystem; |
| Function | interleukin-10 receptor activity; protein binding; receptor activity; signal transducer activity; |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA