

Recombinant Human IL11 protein

Cat. No. IL11-123H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human IL11 protein was expressed in Escherichia coli.
Species	Human
Source	E.coli
ProteinLength	179
Description	Interleukin-11 (IL-11) is encoded by the IL11 gene and is a member of IL-6 superfamily, distinguished based on their use of the common co-receptor gp130. It is a multifunctional cytokine first isolated from bone marrow-derived stromal cells. IL-11 receptor activation requires formation of a complex of two IL-11 molecules with two molecules of the ligand-binding IL-11R α subunit and two molecules of the expressed cell signaling β subunit, gp130. The functions of IL-11 are that it directly stimulates the proliferation of hematopoietic stem cells and megakaryocyte progenitor cells, and induces megakaryocyte maturation resulting in increased platelet production.
Form	Lyophilized from a 0.2 μ m filtered concentrated solution in PBS, pH 7.4.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by a cell proliferation assay using murine B9-11 cells is less than 1 ng/ml, corresponding to a specific activity of > 1.0 \times 10 ⁶ IU/mg.
Molecular Mass	Approximately 19.1 kDa, a single non-glycosylated polypeptide chain containing 179 amino acids.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence	MPGPPPGPPRVSPDPRAELDSTVLLTRSLADTRQLAAQLRDKFPADGDHNLDSLPTLAMSAGALGALQLPGVLTRLRADLLSYLRHVQWLRRAAGSSSLKTLEPELGTQLQARLDRLLRRLQLLMSRLALPQPPDPPAPPLAPPSSAWGGIRAAHAILGGLHLTLDWAVRGLLLLKTRL
Endotoxin	Less than 1 EU/g of rHuIL-11 as determined by LAL method.
Purity	>95% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	IL11
Official Symbol	IL11
Synonyms	IL11; interleukin 11; interleukin-11; adipogenesis inhibitory factor; AGIF; IL 11; oprelvekin; IL-11;
Gene ID	3589

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



mRNA Refseq	NM_000641
Protein Refseq	NP_000632
MIM	147681
UniProt ID	P20809

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA