

Recombinant Human Interleukin 12B

Cat. No. IL12B-177H Lot. No. (See product label)

SPECIFICATION

| | |
|----------------------------|---|
| Product Overview | Recombinant human IL-12 p40 is a 40 kDa protein containing 306 amino acid residues. It was expressed in E.Coli. |
| Species | Human |
| Source | E.coli |
| Description | Interleukin-12 is a disulfide-linked heterodimeric protein (p70), composed of two subunits, p35 and p40, which are encoded by two different genes. Accumulating data indicate that p40 secretion precedes that of IL-12 expression. In addition, to its ability to covalently bind to p35 to form IL-12, p40 can bind to p19 to form IL-23. |
| Purity | > 95% by SDS-PAGE and HPLC analysis. |
| Endotoxin Level | < 0.1 ng per µg of IL-12p40. |
| Stabilizer | None. |
| Formulation | Lyophilized. |
| Biological Activity | Not available. |
| Reconstitution | The human IL-12p40 should be reconstituted in water to a concentration of 0.1-1.0 mg/ml. This solution can be diluted in water or other buffer solutions or stored at -20°C. |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Stability

The lyophilized protein is stable for a few weeks at room temperature, but best stored at -20°C. Reconstituted human IL-12p40 should be stored in working aliquots at -20°C. Avoid repeated freeze-thaw cycles.

GENE INFORMATION

Gene Name

IL12B interleukin 12B (natural killer cell stimulatory factor 2, cytotoxic lymphocyte maturation factor 2, p40) [Homo sapiens]

Synonyms

OTTHUMP00000160820; CLMF; CLMF2; IL-12B; NKSF; NKSF2; CLMF p40; Cytotoxic lymphocyte maturation factor 40 kDa subunit; IL-12 subunit p40; IL12, subunit p40; NK cell stimulatory factor chain 2; cytotoxic lymphocyte maturation factor 2, p40; interleukin 12, p40; interleukin 12B; interleukin 12B (natural killer cell stimulatory factor 2, cytotoxic; interleukin 12B (natural killer cell stimulatory factor 2, cytotoxic lymphocyte maturation factor 2, p40); IL12B

Gene ID

3593

mRNA Refseq

NM_002187

Protein Refseq

NP_002178

MIM

161561

UniProt ID

P29460

Chromosome Location

5q31.1-q33.1

Pathway

Cytokine-cytokine receptor interaction; Jak-STAT signaling pathway; Toll-like receptor signaling pathway; Type I diabetes mellitus

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

contributes_to cytokine activity; hematopoietin/interferon-class (D200-domain) cytokine receptor activity; interleukin-12 alpha subunit binding; interleukin-12 receptor binding; protein binding; protein heterodimerization activity; protein homodimerization activity; signal transducer activity

PDB rendering based on 1f42.



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA