

## Active Recombinant Human IL15RA protein, Fc/Avi-tagged, Biotinylated

**Cat. No.** IL15RA-051H    **Lot. No.** (See product label)

### SPECIFICATION

<b>Product Overview</b>	Biotinylated Recombinant Human IL15RA(Ile31-Thr205) protein, fused to Fc/Avi tag at the C-terminus, was expressed in CHO cells .
<b>Species</b>	Human
<b>Source</b>	CHO
<b>ProteinLength</b>	Ile31-Thr205
<b>Description</b>	Interleukin 15 Receptor alpha (IL-15 R alpha ), also known as CD215, is a widely expressed 60-kDa transmembrane glycoprotein that plays an important role in the homeostasis and activation of NK cells and CD8+ memory T cells and participates in the development and function of many other hematopoietic cell types and non-immune cell types (1-3). Mature human IL-15 R alpha consists of a 175 amino acid (aa) extracellular domain (ECD) containing one N-linked glycosylation site, a 23 aa transmembrane segment, and a 39 aa cytoplasmic tail (4). Within the ECD, human IL-15 R alpha shares approximately 60% aa sequence identity with mouse and rat IL-15 R alpha. Alternate splicing of human IL-15 R alpha generates additional isoforms with variable length deletions in the ECD and/or substitutions in the cytoplasmic domain (4, 5). IL-15 R alpha binds to Interleukin-15 with high affinity (6). IL-15 additionally interacts with lower affinity to a complex of IL-2 R beta and the common gamma chain ( gamma c) which are also subunits of the IL-2 receptor complex (7, 8). The use of shared receptor

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

components contributes to the overlapping biological effects of IL-15 and IL-2. The dominant mechanism of IL-15 action is known as transpresentation in which IL-15/IL-15 R alpha complexes are expressed on the surface of one cell and interact with complexes of IL-2 R beta / gamma c on adjacent cells (9). This enables cells to respond to IL-15 even if they do not express IL-15 R alpha (10-12). IL-15/IL-15 R alpha complexes can transmit reverse signaling that promotes cellular adhesion, tyrosine phosphorylation of intracellular proteins, and cytokine secretion by the IL-15/IL-15 R alpha expressing cells (13, 14). Shed soluble forms of IL-15 R alpha retain the ability to bind tightly to IL-15 and can inhibit IL-15 bioactivity (6, 15, 16). Our Avi-tag Biotinylated Recombinant Human IL-15 R alpha features biotinylation at a single site contained within the Avi-tag, a unique 15 amino acid peptide. Protein orientation will be uniform when bound to streptavidin-coated surface due to the precise control of biotinylation and the rest of the protein is unchanged so there is no interference in the protein's bioactivity.

**Predicted N Terminal** Ile31

**Form** Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose.

**Bio-activity** The biotin to protein ratio is greater than 0.7 as determined by the HABA assay. Measured by its binding ability in a functional ELISA. When Biotinylated Recombinant Human IL15RA Fc Chimera Avi-tag is immobilized at 0.5 µg/mL (100 µ/well), Recombinant Human IL15 binds with an ED50 of 0.02-0.16 ng/mL.

**Molecular Mass** 70-80 kDa, under reducing conditions

**Endotoxin** <0.10 EU per 1 µg of the protein by the LAL method.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

<b>Purity</b>	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
<b>Applications</b>	Binding Activity, Bioactivity
<b>Storage</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 3 months, -20 to -70 °C under sterile conditions after reconstitution.
<b>Reconstitution</b>	Reconstitute at 400 µg/mL in PBS.
<b>Conjugation</b>	Biotin

## GENE INFORMATION

<b>Gene Name</b>	IL15RA interleukin 15 receptor, alpha [ Homo sapiens ]
<b>Official Symbol</b>	IL15RA
<b>Synonyms</b>	IL15RA; interleukin 15 receptor, alpha; interleukin-15 receptor subunit alpha; CD215; IL 15RA; MGC104179;
<b>Gene ID</b>	3601
<b>mRNA Refseq</b>	NM_001243539
<b>Protein Refseq</b>	NP_001230468
<b>MIM</b>	601070

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA




**UniProt ID**

Q13261


**SDS-PAGE**

2 µg/lane Protein was resolved with SDS-PAGE under reducing (R) and non-reducing (NR) conditions and visualized by Coomassie® Blue staining.

**Binding Activity**

 Tel: 1-631-559-9269 1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA