

Active Recombinant Human IL17A protein

Cat. No. IL17A-122H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human IL17A protein(Q16552)(24-155 aa) was expressed in E. coli.
Species	Human
Source	E.coli
ProteinLength	24-155 aa
Form	Lyophilized from a 0.2 µ filtered concentrated solution in 1 × PBS, 5% Trehalose, 0.02% Tween-20.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by inducing IL-6 secretion of murine NIH/3T3 cells is less than 7.5 ng/ml, corresponding to a specific activity of > 1.3 × 10 ⁵ IU/mg.
Molecular Mass	Approximately 31.0 kDa, a disulfide-linked homodimer of two 132 amino acid polypeptide chains.
AASequence	GITIPRNPGC PNPSEDKNFPR TVMVNLNIHN RNTNTNPKRS SDYYNRSTSP WNLHRNEDPE RYPSVIWEAK CRHLGCINAD GNVDYHMNSV PIQQEILVLR REPPHCPNSF RLEKILVSVG CTCVTPIVHH VA
Endotoxin	Less than 1 EU/g of rHuIL-17 as determined by LAL method.
Purity	> 95 % by SDS-PAGE and HPLC analyses.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage

Use a manual defrost freezer and avoid repeated freeze-thaw cycles. - 12 months from date of receipt, -20 to -70 °C as supplied. - 1 month, 2 to 8 °C under sterile conditions after reconstitution. - 3 months, -20 to -70 °C under sterile conditions after reconstitution.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name

[IL17A interleukin 17A \[Homo sapiens \]](#)

Official Symbol

[IL17A](#)

Synonyms

IL17A; interleukin 17A; CTLA8, IL17, interleukin 17 (cytotoxic T lymphocyte associated serine esterase 8); interleukin-17A; cytotoxic T lymphocyte associated protein 8; IL 17; IL 17A; CTLA-8; cytotoxic T-lymphocyte-associated antigen 8; cytotoxic T-lymphocyte-associated protein 8; cytotoxic T-lymphocyte-associated serine esterase 8; interleukin 17 (cytotoxic T-lymphocyte-associated serine esterase 8); IL17; CTLA8; IL-17; IL-17A;

Gene ID

[3605](#)

mRNA Refseq

[NM_002190](#)

Protein Refseq

[NP_002181](#)

MIM

[603149](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



UniProt ID

Q16552

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA