

Active Recombinant Human IL17A Protein, Biotinylated

Cat. No. IL17A-166H **Lot. No.** (See product label)

SPECIFICATION

Product Overview Recombinant Human IL17A(Gly24-Ala155) was expressed in HEK293, Biotinylated.

Species Human

Source HEK293

ProteinLength Gly24-Ala155

Description

Interleukin-17A (IL-17A), also known as CTLA-8, is a 15-20 kDa glycosylated cytokine that plays an important role in anti-microbial and chronic inflammation. The six IL-17 cytokines (IL-17A-F) are encoded by separate genes but adopt a conserved cystine knot fold. Mature human IL-17A shares 60% amino acid sequence identity with mouse and rat IL-17A. IL-17A is secreted by Th17 cells, gamma /delta T cells, iNKT cells, NK cells, LTi cells, neutrophils, and intestinal Paneth cells. It forms disulfide-linked homodimers as well as disulfide-linked heterodimers with IL-17F. IL-17A exerts its effects through the transmembrane IL-17RA in complex with IL-17RC or IL-17RD. Both IL-17RA and IL-17RC are required for responsiveness to heterodimeric IL-17A/F. IL-17A promotes protective mucosal and epidermal inflammation in response to microbial infection. It induces chemokine production, neutrophil influx, and the production of antibacterial peptides. IL-17A/F likewise induces neutrophil migration, but IL-17F does not. IL-17A additionally enhances the production of inflammatory mediators by rheumatoid synovial fibroblasts and contributes to TNF-alpha induced shock. In contrast, it can protect against the progression of colitis by limiting chronic inflammation. IL-17A encourages the formation of autoreactive germinal centers and

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	exacerbates the onset and progression of experimental models of autoimmunity. IL-17A has been shown to exert either tumorigenic or anti-tumor effects (16, 17).
Predicted N Terminal	Gly24
Form	Lyophilized from a 0.2 µm filtered solution in PBS and NaCl with BSA as a carrier protein.
Bio-activity	Measured by its ability to induce IL-6 secretion by NIH-3T3 mouse embryonic fibroblast cells. Yao, Z. et al. (1995) Immunity 3:811. The ED50 for this effect is 1.5-7.5 ng/mL.
Molecular Mass	Predicted Molecular Mass: 15 kDa (unlabeled) SDS-PAGE: 14-23 kDa, reducing conditions
Endotoxin	<0.10 EU per 1 µg of the protein by the LAL method.
Purity	>95%, by SDS-PAGE with silver staining
Notes	Structure / Form: Disulfide-linked homodimer, Biotinylated via sugars
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile
Reconstitution	Reconstitute at 100 µg/mL in PBS containing at least 0.1% human or bovine serum albumin.
Conjugation	Biotin

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

GENE INFORMATION

Gene Name IL17A interleukin 17A [Homo sapiens]

Official Symbol IL17A

Synonyms IL17A; interleukin 17A; CTLA8, IL17, interleukin 17 (cytotoxic T lymphocyte associated serine esterase 8); interleukin-17A; cytotoxic T lymphocyte associated protein 8; IL 17; IL 17A; CTLA-8; cytotoxic T-lymphocyte-associated antigen 8; cytotoxic T-lymphocyte-associated protein 8; cytotoxic T-lymphocyte-associated serine esterase 8; interleukin 17 (cytotoxic T-lymphocyte-associated serine esterase 8); IL17; CTLA8; IL-17; IL-17A;

Gene ID 3605

mRNA Refseq NM_002190

Protein Refseq NP_002181

MIM 603149

UniProt ID Q16552

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA