

Recombinant Human IL17C protein, His-tagged

Cat. No. IL17C-149H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human IL17C(Cys36~Val187) fused with His tag at N-terminal was expressed in E. coli.
Species	Human
Source	E.coli
ProteinLength	Cys36~Val187
Description	The protein encoded by this gene is a T cell-derived cytokine that shares the sequence similarity with IL17. This cytokine was reported to stimulate the release of tumor necrosis factor alpha and interleukin 1 beta from a monocytic cell line. The expression of this cytokine was found to be restricted to activated T cells.
Form	PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300
Molecular Mass	20.7kDa
Endotoxin	<1.0EU per 1ug (determined by the LAL method)
Purity	> 95%
Applications	SDS-PAGE; WB; ELISA; IP
Stability	The thermal stability is described by the loss rate. The loss rate was determined by

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

accelerated thermal degradation test, that is, incubate the protein at 37 centigrade for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Storage

Avoid repeated freeze/thaw cycles. Store at 2-8 centigrade for one month. Aliquot and store at -80 centigrade for 12 months.

Reconstitution

Reconstitute in PBS or others.

GENE INFORMATION

Gene Name

IL17C interleukin 17C [Homo sapiens]

Official Symbol

IL17C

Synonyms

IL17C; interleukin 17C; interleukin-17C; CX2; IL 17C; IL 21; MGC126884; MGC138401; cytokine CX2; IL-21; IL-17C;

Gene ID

27189

mRNA Refseq

NM_013278

Protein Refseq

NP_037410

MIM

604628

UniProt ID

Q9P0M4

**Chromosome
Location**

16q24

Function

cytokine activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA