

Recombinant Human Interleukin 17C Carrier-Free, His-tagged

Cat. No. IL17C-293H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human Carrier-Free IL-17C with His-tag was expressed in <i>Insect cell</i> . MW= 19.793 kDa.
Species	Human
Source	Insect Cells
Description	IL-17C is a member of the IL-17 family, which is comprised of six proteins (IL-17A through IL-17F). IL-17 proteins are pro-inflammatory cytokines that induce local cytokine production and are involved in the regulation of immune functions. IL-17C has been reported to stimulate the release of TNF-alpha and IL-1 β from the monocytic cell line THP-1. IL-17C shares 15% to 30% amino acid sequence identity with the other IL-17 family members. Mouse and human IL-17C share 83% amino acid sequence identity. Human IL-17C cDNA encodes a 197 amino acid protein with a putative 18 amino acid signal peptide. The recombinant human IL-17C is a disulfide-linked homodimer composed of a 19.8 kDa monomeric subunit.
Handling Conditions	For best recovery, quick-spin vial prior to opening. Use in a sterile environment.
Purity	Greater than 98% as determined by SDS-PAGE.
Endotoxin Level	Less than 0.01 ng/ug cytokine as determined by the LAL assay.
Bioactivity	Testing in progress.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Formulation 20 mM Phosphate, 0.2 M NaCl, pH 6.0, with 1.0% BSA.

Temperature Limitation Store at less than or equal to -70°C.

GENE INFORMATION

Gene Name IL17C interleukin 17C [Homo sapiens]

Synonyms interleukin 17C; CX2; IL-21; IL-17C; MGC126884; MGC138401; IL17C; interleukin-17C; cytokine CX2; Cytokine CX2

Gene ID 27189

mRNA Refseq NM_013278

Protein Refseq NP_037410

UniProt ID Q9P0M4

Chromosome Location 16q24

MIM 604628

Function cytokine activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA