

Active Recombinant Human IL17F Protein (133 aa)

Cat. No. IL17F-099I Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human IL17F Protein without tag was expressed in E. coli.
Species	Human
Source	E.coli
ProteinLength	133
Description	Human IL-17F is synthesized as a 153 aa precursor with a 20 aa signal sequence and a 133 aa mature region. Like IL-17A, IL-17F contains one potential site for N-linked glycosylation. IL-17A and IL-17F share 50% aa sequence identity. IL17-F homodimer is produced by an activated subset of CD4+ T cells, termed Th17. IL17-F has been shown to stimulate proliferation and activation of T-cells and PBMCs. IL-17F also regulates cartilage matrix turnover and inhibits angiogenesis.
Form	Sterile Filtered White lyophilized (freeze-dried) powder.
Bio-activity	Measured by its ability to induce IL-6 production by NHDF cells, it's fully biologically active when compared to standard.
Molecular Mass	A disulfide-linked homodimer of 30.1kDa, consisting of two 133 amino acid polypeptide chains.
AA Sequence	MRKIPKVGHTFFQKPESCPPVPGGSMKLDIGIINENQRVSMRNIESRSTSPWNYTV TWDPNRYPSEVVQAQCRNLGCINAQGKEDISMNSVPIQQETLVVRRKHQGCVSF

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

QLEKVLVTVGCTCVTPVIHHVQ

Endotoxin Less than 1 EU/μg of rHuIL-17F as determined by LAL method.

Purity >95% by SDS-PAGE and HPLC analyses.

Storage This lyophilized preparation is stable for several weeks at 2-8 centigrade, but should be kept at -20 centigrade for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 centigrade. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 to -70 centigrade. Avoid repeated freeze/thaw cycles.

Storage Buffer Lyophilized from a 0.2 μm filtered concentrated solution in PBS, pH 7.4.

Reconstitution We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in 4 mM HCl to a concentration of 0.1mg/ml. Stock solutions should be apportioned into working aliquots and stored at < -20C. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name [IL17F interleukin 17F \[Homo sapiens \]](#)

Official Symbol [IL17F](#)

Synonyms IL17F; interleukin 17F; interleukin-17F; IL 17F; ML 1; ML1; IL-24; cytokine ML-1; mutant IL-17F; interleukin-24; ML-1; CANDF6; IL-17F;

Gene ID [112744](#)

mRNA Refseq [NM_052872](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Protein Refseq NP_443104

MIM 606496

UniProt ID Q96PD4

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA