

Active Recombinant Rhesus IL17RA Protein, His-tagged

Cat. No. IL17RA-442R Lot. No. (See product label)

SPECIFICATION

| | |
|-------------------------|--|
| Product Overview | Recombinant Rhesus IL17RA (XP_001102483.2) (Met1-Trp320) was expressed with a polyhistidine tag at the C-terminus. |
| Species | Rhesus macaque |
| Source | HEK293 |
| ProteinLength | Met1-Trp320 |
| Description | <p>Interleukin-17 receptor (IL-17R), also known as Interleukin-17 receptor A (IL-17RA) and CD217 antigen (CD217), is a cytokine receptor which binds interleukin 17. IL-17R/IL-17RA (CD217) is a proinflammatory cytokine secreted by activated T-lymphocytes. It is a potent inducer of the maturation of CD34-positive hematopoietic precursors into neutrophils. IL-17R/IL-17RA (CD217) is a ubiquitous type I membrane glycoprotein that binds with low affinity to interleukin 17A. Interleukin 17A and its receptor IL-17RA play a pathogenic role in many inflammatory and autoimmune diseases such as rheumatoid arthritis. Like other cytokine receptors, this receptor likely has a multimeric structure. Defects in IL-17R/IL-17RA (CD217) are the cause of familial candidiasis type 5 (CANDF5). CANDF5 is a rare disorder with altered immune responses and impaired clearance of fungal infections, selective against <i>Candida</i>. It is characterized by persistent and/or recurrent infections of the skin, nails and mucous membranes caused by organisms of the genus <i>Candida</i>, mainly <i>Candida albicans</i>.</p> |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

| | |
|-----------------------------|---|
| Predicted N Terminal | Leu 33 |
| Form | Lyophilized from sterile PBS, pH 7.4. Normally 5 %- 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. |
| Bio-activity | Measured by its binding ability in a functional ELISA. Immobilized rhesus IL17RA-His at 10µg/mL (100µL/well) can bind biotinylated human IL17A, the EC50 of biotinylated human IL17A is 15-100ng/mL. |
| Molecular Mass | The recombinant rhesus IL17RA consists of 309 amino acids and predicts a molecular mass of 35.6 kDa |
| Endotoxin | < 1.0 EU per µg protein as determined by the LAL method. |
| Purity | > 95 % as determined by SDS-PAGE. |
| Storage | Samples are stable for up to twelve months from date of receipt at -70 centigrade. Store it under sterile conditions at -20 centigrade to -80 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. |
| Reconstitution | It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents. |

GENE INFORMATION

| | |
|------------------------|---------------------------------------|
| Official Symbol | IL17RA |
| Synonyms | IL17A;interleukin 17A;interleukin-17A |

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA